

EXHIBIT 1

(12) **United States Patent**
Wilton et al.

(10) **Patent No.:** **US 8,455,636 B2**
(45) **Date of Patent:** **Jun. 4, 2013**

(54) **ANTISENSE OLIGONUCLEOTIDES FOR INDUCING EXON SKIPPING AND METHODS OF USE THEREOF**

(75) Inventors: **Stephen Donald Wilton**, Applecross (AU); **Sue Fletcher**, Bayswater (AU); **Graham McClorey**, Bayswater (AU)

(73) Assignee: **The University of Western Australia**, Crawley (AU)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **13/271,080**

(22) Filed: **Oct. 11, 2011**

(65) **Prior Publication Data**

US 2012/0022145 A1 Jan. 26, 2012

Related U.S. Application Data

(63) Continuation of application No. 12/837,359, filed on Jul. 15, 2010, now Pat. No. 8,232,384, which is a continuation of application No. 11/570,691, filed as application No. PCT/AU2005/000943 on Jun. 28, 2005, now Pat. No. 7,807,816.

(30) **Foreign Application Priority Data**

Jun. 28, 2004 (AU) 2004903474

(51) **Int. Cl.**
C07H 21/04 (2006.01)

(52) **U.S. Cl.**
USPC **536/24.5**; 536/24.31; 536/24.1; 514/44

(58) **Field of Classification Search**
None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,034,506 A 7/1991 Summerton et al.
5,142,047 A 8/1992 Summerton et al.
5,149,797 A 9/1992 Pederson et al.
5,166,315 A 11/1992 Summerton et al.
5,185,444 A 2/1993 Summerton et al.
5,217,866 A 6/1993 Summerton et al.
5,506,337 A 4/1996 Summerton et al.
5,521,063 A 5/1996 Summerton et al.
5,627,274 A 5/1997 Kole et al.
5,665,593 A 9/1997 Kole et al.
5,869,252 A 2/1999 Bouma et al.
5,892,023 A 4/1999 Pirotzky et al.
6,153,436 A 11/2000 Hermonat et al.
6,210,892 B1 4/2001 Bennett et al.
6,653,466 B2 11/2003 Matsuo
6,653,467 B1 11/2003 Matsuo et al.
6,656,732 B1 12/2003 Bennett et al.
6,727,355 B2 4/2004 Matsuo et al.
6,784,291 B2 8/2004 Iversen et al.
7,070,807 B2 7/2006 Mixson
7,163,695 B2 1/2007 Mixson
7,250,289 B2 7/2007 Zhou
7,314,750 B2 1/2008 Zhou

7,468,418 B2 12/2008 Iversen et al.
7,534,879 B2 * 5/2009 van Deutekom 536/24.5
7,655,785 B1 2/2010 Bentwich
7,807,816 B2 10/2010 Wilton et al.
7,902,160 B2 3/2011 Matsuo et al.
7,960,541 B2 6/2011 Wilton et al.
7,973,015 B2 7/2011 van Ommen et al.
8,084,601 B2 12/2011 Popplewell et al.
8,324,371 B2 12/2012 Popplewell et al.
2002/0156235 A1 * 10/2002 Manoharan et al. 530/322
2003/0224353 A1 12/2003 Stein et al.
2004/0248833 A1 12/2004 Emanuele et al.
2004/0254137 A1 * 12/2004 Ackermann et al. 514/44
2005/0026164 A1 2/2005 Zhou
2005/0153935 A1 7/2005 Iversen et al.
2006/0099616 A1 5/2006 van Ommen et al.
2006/0147952 A1 7/2006 van Ommen et al.
2007/0082861 A1 4/2007 Matsuo et al.
2008/0209581 A1 8/2008 van Ommen et al.
2009/0228998 A1 9/2009 van Ommen et al.
2009/0269755 A1 10/2009 Aartsma-Rus et al.
2009/0312532 A1 12/2009 Van Deutekom et al.
2010/0130591 A1 5/2010 Sazani et al.
2011/0015253 A1 1/2011 Wilton et al.
2011/0015258 A1 1/2011 Wilton et al.
2011/0046203 A1 2/2011 Wilton et al.
2011/0046360 A1 2/2011 Matsuo et al.
2011/0263682 A1 10/2011 De Kimpe et al.
2011/0294753 A1 12/2011 De Kimpe et al.

(Continued)

FOREIGN PATENT DOCUMENTS

AU 780517 11/2001
AU 2003284638 A1 6/2004

(Continued)

OTHER PUBLICATIONS

Aartsma-Rus et al., "Antisense-Induced Multiexon Skipping for Duchenne Muscular Dystrophy Makes More Sense," *Am. J. Hum. Genet.* 74:83-92, 2004.
Aartsma-Rus et al., "Targeted exon skipping as a potential gene correction therapy for Duchenne muscular dystrophy," *Neuromuscular Disorders* 12:S71-S77, 2002.
Aartsma-Rus et al., "Therapeutic antisense-induced exon skipping in cultured muscle cells from six different DMD patients," *Human Molecular Genetics* 12(8):907-914, 2003.
Abbs et al., "A convenient multiplex PCR system for the detection of dystrophin gene deletions: a comparative analysis with cDNA hybridisation shows mistyping by both methods," *J. Med. Genet.* 28:304-311, 1991.
Agrawal et al., "Oligodeoxynucleoside phosphoramidates and phosphorothioates as inhibitors of human immunodeficiency virus," *Proc. Natl. Acad. Sci. USA* 85:7079-7083, Oct. 1988.
Akhtar et al., "Cellular uptake and intracellular fate of antisense oligonucleotides," *Trends in Cell Biology* 2:139-144, May 1992.
Akhtar, Saghir, (ed), *Delivery Strategies for antisense Oligonucleotide Therapeutics*, CRC Press Boca Raton, Florida, 1995.

(Continued)

Primary Examiner — Kimberly Chong

(74) *Attorney, Agent, or Firm* — Nelson Mullins Riley & Scarborough LLP; Amy E. Mandragouras, Esq.; Erika L. Wallace

(57) **ABSTRACT**

An antisense molecule capable of binding to a selected target site to induce exon skipping in the dystrophin gene, as set forth in SEQ ID NO: 1 to 202.

43 Claims, 22 Drawing Sheets

US 8,455,636 B2

Page 2

U.S. PATENT DOCUMENTS

2011/0312086	A1	12/2011	Van Deutekom
2012/0022134	A1	1/2012	De Kimpe et al.
2012/0108652	A1	5/2012	Popplewell et al.
2012/0108653	A1	5/2012	Popplewell et al.
2012/0172415	A1	7/2012	Voit et al.

FOREIGN PATENT DOCUMENTS

CA	2 507 125	A1	6/2004
EP	1054058	A1	11/2000
EP	1160318	A2	12/2001
EP	1 191 097	A1	3/2002
EP	1191098	A2	3/2002
EP	1544297	A2	6/2005
EP	1568769	A1	8/2005
EP	1619249	A1	1/2006
EP	1 766 010	AO	3/2007
EP	1857548	A1	11/2007
EP	2135948	A2	12/2009
EP	2284264	A1	2/2011
EP	2374885	A2	10/2011
EP	2386636	A2	11/2011
EP	2392660	A2	12/2011
EP	2530153	A1	12/2012
EP	2530154	A1	12/2012
EP	2530155	A1	12/2012
EP	2530156	A1	12/2012
WO	93/20227	A1	10/1993
WO	WO 94/02595	A1	2/1994
WO	WO 96/10391	A1	4/1996
WO	WO 96/10392	A1	4/1996
WO	WO 97/30067	A1	8/1997
WO	WO 97/34638	A1	9/1997
WO	WO 00/44897	A1	8/2000
WO	WO 01/49775	A2	7/2001
WO	WO 01/83740	A2	11/2001
WO	WO 02/024906	A1	3/2002
WO	03/053341	A2	7/2003
WO	WO 2004/048570	A1	6/2004
WO	WO 2004/083432	A1	9/2004
WO	WO 2004/083446	A2	9/2004
WO	WO 2006/000057	A1	1/2006
WO	2006/112705	A2	10/2006
WO	2007/135105	A1	11/2007
WO	2009/054725	A2	4/2009
WO	2009/101399	A1	8/2009
WO	2009/139630	A2	11/2009
WO	2010/050801	A1	5/2010
WO	2010/050802	A2	5/2010
WO	2010/115993	A1	10/2010
WO	2010/123369	A1	10/2010
WO	2010/150231	A1	12/2010
WO	2011/024077	A2	3/2011
WO	2011/057350	A1	5/2011
WO	2012/001941	A1	1/2012
WO	2012/029986	A1	3/2012
WO	2012/109296	A1	8/2012

OTHER PUBLICATIONS

Anderson, "Human Gene Therapy," *Science* 256:808-813, May 8, 1992.

Asvadi et al., "Expression and functional analysis of recombinant scFv and diabody fragments with specificity for human RhD," *Journal of Molecular Recognition* 15:321-330, 2002.

Berge et al., "Pharmaceutical Salts," *Journal of Pharmaceutical Sciences* 66(1):1-19, Jan. 1977.

Brown et al., "Dystrophic phenotype induced in vitro by antibody blockade of muscle ctdystroglycan-laminin interaction," *Journal of Cell Science* 112:209-216, 1999.

Collins et al., "Duchenne's muscular dystrophy: animal models used to investigate pathogenesis and develop therapeutic strategies," *International Journal of Experimental Pathology* 84:165-172, 2003. *Journal of Experimental Pathology* 84:163-172, 2003.

De Angelis et al., "Chimeric snRNA molecules carrying antisense sequences against the splice junctions of exon 51 of the dystrophin pre-mRNA induce exon skipping and restoration of a dystrophin synthesis in 448-50 DMD cells," *Proc. Natl. Acad. Sci. USA* 99(14):9456-9461, Jul. 9, 2002.

DelloRusso et al., "Functiontd correction of adult *mdx* mouse muscle using gutted adenoviral vectors expressing full-length dystrophin," *Proc. Natl. Acad. Sci. USA* 99(20):12979-12984, Oct. 2002.

Dirksen et al., "Mapping the SF2/ASF Binding Sites in the Bovine Growth Hormone Exonic Splicing Enhancer," *Journal of Biological Chemistry* 275(37):29170-29177, 2000.

Dunckley et al., "Modulation of Splicing in the dystrophine gene in cultures *Mdx* muscle cells by antisense oligoribuncleotides," *Human Molecular Genetics* 5(1):1083-1090, 1995.

Dunckley et al., "Modulation of splicing in the *DMD* Gene By Antisense Oligoribuncleotides," *Nucleosides & Nucleotides* 16(7-9):1665-1668, 1997.

Errington et al., "Target selection for antisense oligonucleotide induced exon skipping in the dystrophin gene," *The Journal of Gene Medicine* 5:518-527, 2003.

"Exon 51 Sequence of Dystrophin," Document D19 as filed in Opposition of European Patent EP1619249, filed Jun. 23, 2009, 7 pages.

Friedmann, "Progress Toward Human Gene Therapy," *Science* 244:1275-1281, Jun. 16, 1989.

Gebski et al., "Moipholino antisense oligonucleotide induced dystrophin exon 23 skipping in *mdx* mouse muscle," *Human Molecular Genetics* 12(15):1801-1811, 2003.

Gennaro, Alfonso R., *Remington's Pharmaceutical Studies, 18th Edition*, Mack Publishing, Co., Eastern, PA. 1990.

Giles et al., "Antisense Morpholino Oligonucleotide Analog Induces Missplicing of *C-myc* mRNA," *Antisense & Nucleic Acid Drug Development* 9:213-220, 1999.

Harel-Bellan et al., "Specific Inhibition of *c-myc* Protein Biosynthesis Using An Antisense Synthetic Deoxy-Oligonucleotide In Human T Lymphocytes," *The Journal of Immunology* 140(7):2431-2435, Apr. 1, 1988.

Hussey et al., "Analysis of five Duchenne muscular dystrophy exons and gender determination using conventional duplex polymerase chain reaction on single cells," *Molecular Human Reproduction* 5(11):1089-1094, 1999.

Karras et al., "Deletion of Individual Exons and Induction of Soluble Murine Interleukin-5 Receptor- α Chain Expression through Antisense Oligonucleotide-Mediated Redirection of PremRNA Splicing," *Molecular Pharmacology* 58:380-387, 2000.

Liu et al., "Identification of functional exonic splicing enhancer motifs recognized by individual SR proteins," *Genes & Development* 12:1998-2012, 1998.

Lu et al., "Functional amounts of dystrophin produced by skipping the mutated exon in the *mdx* dystrophic mouse," *Nature Medicine* 9(8):1009-1014, Aug. 2003.

Lu et al., "Massive idiosyncratic Exon Skipping Corrects the Non-sense Mutation in Dystrophic Mouse Muscle and Produces Functional Revertant Fibers by Clonal Expansion," *The Journal of Cell Biology* 148(5):985-995, Mar. 6, 2000.

Mann et al., "Antisense-induced exon skipping and synthesis of dystrophin in the *mdx* mouse," *Proc. Natl. Acad. Sci. USA* 98(1):42-47, Jan. 2, 2001.

Mann et al., "Improved antisense oligonucleotide induced exon skipping in the *mdx* mouse model of muscular dystrophy," *The Journal of Gene Medicine* 4:644-654, 2002.

Matsuo et al., "Exon Skipping during Splicing of Dystrophin mRNA Precursor due to an intraexon Deletion in the Dystrophin Gene of Duchenne Muscular Dystrophy Kobe," *J. Clin. Invest.* 87:2127-2131, Jun. 1991.

Matsuo, "Duchenne and Becker Muscular Dystrophy: From Gene Diagnosis to Molecular Therapy," *IUBMB Life* 53:147-152, 2002.

Matsuo, "Duchenne/Becker muscular dystrophy: from molecular diagnosis to gene therapy," *Brain & Development* 18:167-172, 1996.

Monaco et al., "An Explanation for the Phenotypic Differences between Patients Bearing Partial Deletions of the DMD Locus," *Genomics* 2:90-95, 1988.

Pramono et al., "Induction of Exon Skipping of the Dystrophin Transcript in Lymphoblastoid Cells by Transfecting an Antisense Oligodeoxynucleotide Complementary to an Exon Recognition Sequence," *Biochemical and Biophysical Research Communications* 226:445-449, 1996.

Roberts et al., "Exon Structure of the Human Dystrophin Gene," *Genomics* 16:536-538, 1993.

US 8,455,636 B2

Page 3

- Rosso et al., "An *Arabidopsis thaliana* T-DNA mutagenized population (GABI-Kat) for flanking sequence tag-based reverse genetics," *Plant Molecular Biology* 53:247-259, 2003.
- Shapiro et al., "RNA splice junctions of different classes of eukaryotes: sequence statistics and functional implications in gene expression," *Nucleic Acids Research* 15(17):7155-7174, 1987.
- Sherratt et al., "Exon Skipping and Translation in Patients with Frameshift Deletions in the Dystrophin Gene," *Am. J. Hum. Genet.* 53:1007-1015, 1993.
- Shiga et al., "Disruption of the Splicing Enhancer Sequence within Exon 27 of the Dystrophin Gene by a Nonsense Mutation Induces Partial Skipping of the Exon and Is Responsible for Becker Muscular Dystrophy," *J. Clin. Invest.* 100(9):2204-2210, Nov. 1997.
- Sierakowska et al., "Repair of thalassemic human α -globin mRNA in mammalian cells by antisense oligonucleotides," *Proc. Natl. Acad. Sci. USA* 93:12840-12844, Nov. 1996.
- Summerton et al., "Morpholino Antisense Oligomers: Design, Preparation, and Properties," *Antisense & Nucleic Acid Drug Development* 7:187-195, 1997.
- Takeshima et al., "Modulation of In Vitro Splicing of the Upstream Intron by Modifying an IntraExon Sequence Which is Deleted from the Dystrophin Gene in Dystrophin Kobe," *J. Clin. Invest.* 95:515-520, Feb. 1995.
- Tanaka et al., "Polypurine Sequences within a Downstream Exon Function as a Splicing Enhancer," *Molecular and Cellular Biology* 14(2):1347-1354, Feb. 1994.
- Thanh et al., "Characterization of Revertant Muscle Fibers in Duchenne Muscular Dystrophy, Using Exon-Specific Monoclonal Antibodies Against Dystrophin," *Am. J. Hum. Genet.* 56:725-731, 1995.
- van Deutekom et al., "Advances in Duchenne Muscular Dystrophy Gene Therapy," *Nature Reviews Genetics* 4(10):774-783, Oct. 2003.
- van Deutekom et al., "Antisense-induced exon skipping restores dystrophin expression in DMD patient derived muscle cells," *Human Molecular Genetics* 10(15):1547-1554, 2001.
- Volloch et al. "Inhibition of Pre-mRNA Splicing by Antisense RNA In Vitro: Effect of RNA Containing Sequences Complementary To Exons," *Biochemical and Biophysical Research Communications* 179(3):1593-1599, Sep. 30, 1991.
- Watakabe et al., "The role of exon sequences in splice site selection," *Genes & Development* 7:407-418, 1993.
- Wilton et al., "Specific removal of the nonsense mutation from the mdx dystrophin mRNA using antisense oligonucleotides," *Neuromuscular Disorders* 9:330-338, 1999.
- PCT/US01/14410 International Search Report mailed Mar. 6, 2002, (5 pages).
- Patentee's Response to European Patent Application No. 05076770. 6, dated Jul. 28, 2006, 4 pages.
- Errington, Stephen J. et al., "Target selection for antisense oligonucleotide induced exon skipping in the dystrophin gene," *The Journal of Gene Medicine*, vol. 5:518-527 (2003).
- Fletcher, Susan et al., "Gene therapy and molecular approaches to the treatment of hereditary muscular disorders," *Curr. Opin. Neurol.*, vol. 13:553-560 (2000).
- McCloy, Graham et al., "Splicing intervention for Duchenne muscular dystrophy," *Current Opinion in Pharmacology*, vol. 5:529-534 (2005).
- Wilton, Stephen D. et al., "Antisense oligonucleotides in the treatment of Duchenne muscular dystrophy: where are we now?" *Neuromuscular Disorders*, vol. 15:399-402 (2005).
- Cirak, Sebahattin et al., "Exon skipping and dystrophin restoration in patients with Duchenne muscular dystrophy after systemic phosphorodiamidate morpholino oligomer treatment: an open-label, phase 2, dose-escalation study," *Lancet*, vol. 378(9791):595-605 (2011).
- Dominski, Zbigniew et al., "Identification and Characterization by Antisense Oligonucleotides of Exon and Intron Sequences Required for Splicing," *Molecular and Cellular Biology*, vol. 14(11):7445-7454 (1994).
- Dominski, Zbigniew et al., "Restoration of correct splicing in thalassemic pre-mRNA by antisense oligonucleotides," *Proc. Natl. Acad. Sci. USA*, vol. 90:8673-8677 (1993).
- Fall, Abbie M. et al., "Induction of revertant fibres in the mdx mouse using antisense oligonucleotides," *Genetics Vaccines and Therapy*, vol. 4:3, doi:10.1186/1479-0556-4-3, 12 pages (2006).
- Fletcher, Susan et al., "Dystrophin expression in the mdx mouse after localised and systemic administration of a morpholino antisense oligonucleotide," *J. Gene Med.*, vol. 8:207-216 (2006).
- Kaye, Ed, "Results of the Eteplirsen Phase 2b and Phase 2b Extension Study in Duchenne Muscular Dystrophy," 8th Annual Meeting of the Oligonucleotide Therapeutics Society, Session 9: Advances in Oligonucleotide Clinical Development II, p. 48 (2012).
- Kinali, Maria et al., "Local restoration of dystrophin expression with the morpholino oligomer AVI-4658 in Duchenne muscular dystrophy: a single-blind, placebo-controlled, dose-escalation, proof-of-concept study," *Lancet Neurol.*, vol. 8:918-928 (2009).
- Matsuo, Masafumi et al., "Treatment of Duchenne Muscular Dystrophy with Oligonucleotides against an Exonic Splicing Enhancer Sequence," *Basic Appl. Myol.*, vol. 13(6):281-285 (2003).
- McCloy, G. et al., "Antisense oligonucleotide-induced exon skipping restores dystrophin expression in vitro in a canine model of DMD," *Gene Therapy*, vol. 13:1373-1381 (2006).
- McCloy, G. et al., "Induced dystrophin exon skipping in human muscle explants," *Neuromuscular Disorders*, vol. 16:583-590 (2006).
- Mitrant, Chalermchai et al., "Rational Design of Antisense Oligomers to Induce Dystrophin Exon Skipping," *Molecular Therapy*, vol. 17(8):1418-1426 (2009).
- European Search Report for Application No. 10004274.6, 12 pages, dated Jan. 2, 2013.
- Partial European Search Report for Application No. 10004274.6, 6 pages, dated Oct. 2, 2012.
- Partial European Search Report for Application No. 12162995.0, 6 pages, dated Oct. 2, 2012.
- European Search Report for Application No. 12162995.0, 11 pages, dated Jan. 15, 2013.

* cited by examiner

U.S. Patent

Jun. 4, 2013

Sheet 1 of 22

US 8,455,636 B2

FIGURE 1



U.S. Patent

Jun. 4, 2013

Sheet 2 of 22

US 8,455,636 B2




FIGURE 2

U.S. Patent

Jun. 4, 2013

Sheet 3 of 22

US 8,455,636 B2

H8A(-06+14) | H8A(-06+18)
M 600 300 100 50 20 UT 600 300 100 50 20 UT M




FIGURE 3

U.S. Patent

Jun. 4, 2013

Sheet 4 of 22

US 8,455,636 B2

H7A(+45+67) H7A(+2+26)
M 600 300 100 50 20 600N M 600 300 100 50 20 600N M




FIGURE 4

U.S. Patent

Jun. 4, 2013

Sheet 5 of 22

US 8,455,636 B2




FIGURE 5

U.S. Patent

Jun. 4, 2013

Sheet 6 of 22

US 8,455,636 B2

6A(+69+91)

M 600 300 200 100 50 20 UT




FIGURE 6

U.S. Patent

Jun. 4, 2013

Sheet 7 of 22

US 8,455,636 B2

H4A(+13+32)

M 600 300 100 50 20 UT Neg M




FIGURE 7

U.S. Patent

Jun. 4, 2013

Sheet 8 of 22

US 8,455,636 B2




FIGURE 8

U.S. Patent

Jun. 4, 2013

Sheet 9 of 22

US 8,455,636 B2




FIGURE 9

U.S. Patent

Jun. 4, 2013

Sheet 10 of 22

US 8,455,636 B2




FIGURE 10

U.S. Patent

Jun. 4, 2013

Sheet 11 of 22

US 8,455,636 B2




FIGURE 11

U.S. Patent

Jun. 4, 2013

Sheet 12 of 22

US 8,455,636 B2




FIGURE 12

U.S. Patent

Jun. 4, 2013

Sheet 13 of 22

US 8,455,636 B2




FIGURE 13

U.S. Patent

Jun. 4, 2013

Sheet 14 of 22

US 8,455,636 B2




FIGURE 14

U.S. Patent

Jun. 4, 2013

Sheet 15 of 22

US 8,455,636 B2




FIGURE 15

U.S. Patent

Jun. 4, 2013

Sheet 16 of 22

US 8,455,636 B2




FIGURE 16

U.S. Patent

Jun. 4, 2013

Sheet 17 of 22

US 8,455,636 B2





FIGURE 17

U.S. Patent

Jun. 4, 2013

Sheet 18 of 22

US 8,455,636 B2



U.S. Patent

Jun. 4, 2013

Sheet 19 of 22

US 8,455,636 B2

H40A(-05+17)




FIGURE 19

U.S. Patent

Jun. 4, 2013

Sheet 20 of 22

US 8,455,636 B2




FIGURE 20

U.S. Patent

Jun. 4, 2013

Sheet 21 of 22

US 8,455,636 B2

H46A(+86+115)

600 300 200 100 L2K UT




FIGURE 21

U.S. Patent

Jun. 4, 2013

Sheet 22 of 22

US 8,455,636 B2




FIGURE 22

US 8,455,636 B2

1

ANTISENSE OLIGONUCLEOTIDES FOR INDUCING EXON SKIPPING AND METHODS OF USE THEREOF

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a continuation of U.S. patent application Ser. No. 12/837,359, filed Jul. 15, 2010, which is a continuation of U.S. patent application Ser. No. 11/570,691, filed Jan. 15, 2008, now U.S. Pat. No. 7,807,816, which is a 35 U.S.C. §371 National Phase Application of PCT/AU2005/000943, filed Jun. 28, 2005, which claims priority to Australian Patent Application No. 2004903474, filed Jun. 28, 2004; these applications are incorporated herein by reference in their entireties.

STATEMENT REGARDING SEQUENCE LISTING

The instant application contains a Sequence Listing which has been submitted in ASCII format via EFS-Web and is hereby incorporated by reference in its entirety. Said ASCII copy, created on Jan. 30, 2013, is named SequenceListing.txt and is 61 Kilobytes in size. The Sequence Listing is being submitted by EFS Web and is hereby incorporated by reference into the specification.

FIELD OF THE INVENTION

The present invention relates to novel antisense compounds and compositions suitable for facilitating exon skipping. It also provides methods for inducing exon skipping using the novel antisense compounds as well as therapeutic compositions adapted for use in the methods of the invention.

BACKGROUND ART

Significant effort is currently being expended researching methods for suppressing or compensating for disease-causing mutations in genes. Antisense technologies are being developed using a range of chemistries to affect gene expression at a variety of different levels (transcription, splicing, stability, translation). Much of that research has focused on the use of antisense compounds to correct or compensate for abnormal or disease-associated genes in a myriad of different conditions.

Antisense molecules are able to inhibit gene expression with exquisite specificity and because of this many research efforts concerning oligonucleotides as modulators of gene expression have focused on inhibiting the expression of targeted genes such as oncogenes or viral genes. The antisense oligonucleotides are directed either against RNA (sense strand) or against DNA where they form triplex structures inhibiting transcription by RNA polymerase II. To achieve a desired effect in specific gene down-regulation, the oligonucleotides must either promote the decay of the targeted mRNA or block translation of that mRNA, thereby effectively preventing de novo synthesis of the undesirable target protein.

Such techniques are not useful where the object is to up-regulate production of the native protein or compensate for mutations which induce premature termination of translation such as nonsense or frame-shifting mutations. Furthermore, in cases where a normally functional protein is prematurely terminated because of mutations therein, a means for restoring some functional protein production through antisense

2

technology has been shown to be possible through intervention during the splicing processes (Sierakowska H, et al., (1996) *Proc Natl Acad Sci USA* 93, 12840-12844; Wilton S D, et al., (1999) *Neuromusc Disorders* 9, 330-338; van Deutekom J C et al., (2001) *Human Mol Genet.* 10, 1547-1554). In these cases, the defective gene transcript should not be subjected to targeted degradation so the antisense oligonucleotide chemistry should not promote target mRNA decay.

In a variety of genetic diseases, the effects of mutations on the eventual expression of a gene can be modulated through a process of targeted exon skipping during the splicing process. The splicing process is directed by complex multi-particle machinery that brings adjacent exon-intron junctions in pre-mRNA into close proximity and performs cleavage of phosphodiester bonds at the ends of the introns with their subsequent reformation between exons that are to be spliced together. This complex and highly precise process is mediated by sequence motifs in the pre-mRNA that are relatively short semi-conserved RNA segments to which bind the various nuclear splicing factors that are then involved in the splicing reactions. By changing the way the splicing machinery reads or recognises the motifs involved in pre-mRNA processing, it is possible to create differentially spliced mRNA molecules. It has now been recognised that the majority of human genes are alternatively spliced during normal gene expression, although the mechanisms invoked have not been identified. Using antisense oligonucleotides, it has been shown that errors and deficiencies in a coded mRNA could be bypassed or removed from the mature gene transcripts.

In nature, the extent of genetic deletion or exon skipping in the splicing process is not fully understood, although many instances have been documented to occur, generally at very low levels (Sherrat T G, et al., (1993) *Am J Hum Genet.* 53, 1007-1015). However, it is recognised that if exons associated with disease-causing mutations can be specifically deleted from some genes, a shortened protein product can sometimes be produced that has similar biological properties of the native protein or has sufficient biological activity to ameliorate the disease caused by mutations associated with the target exon (Lu Q L, et al., (2003) *Nature Medicine* 9, 1009-1014; Aartsma-Rus A et al., (2004) *Am J Hum Genet.* 74: 83-92).

This process of targeted exon skipping is likely to be particularly useful in long genes where there are many exons and introns, where there is redundancy in the genetic constitution of the exons or where a protein is able to function without one or more particular exons (e.g. with the dystrophin gene, which consists of 79 exons; or possibly some collagen genes which encode for repeated blocks of sequence or the huge nebulin or titin genes which are comprised of ~80 and over 370 exons, respectively).

Efforts to redirect gene processing for the treatment of genetic diseases associated with truncations caused by mutations in various genes have focused on the use of antisense oligonucleotides that either: (1) fully or partially overlap with the elements involved in the splicing process; or (2) bind to the pre-mRNA at a position sufficiently close to the element to disrupt the binding and function of the splicing factors that would normally mediate a particular splicing reaction which occurs at that element (e.g., binds to the pre-mRNA at a position within 3, 6, or 9 nucleotides of the element to be blocked).

For example, modulation of mutant dystrophin pre-mRNA splicing with antisense oligoribonucleotides has been reported both in vitro and in vivo. In one type of dystrophin mutation reported in Japan, a 52-base pair deletion mutation causes exon 19 to be removed with the flanking introns during

US 8,455,636 B2

3

the splicing process (Matsuo et al., (1991) *J Clin Invest.* 87:2127-2131). An in vitro minigene splicing system has been used to show that a 31-mer 2'-O-methyl oligoribonucleotide complementary to the 5' half of the deleted sequence in dystrophin Kobe exon 19 inhibited splicing of wild-type pre-mRNA (Takeshima et al. (1995), *J. Clin. Invest.*, 95, 515-520). The same oligonucleotide was used to induce exon skipping from the native dystrophin gene transcript in human cultured lymphoblastoid cells.

Dunckley et al., (1997) *Nucleosides & Nucleotides*, 16, 1665-1668 described in vitro constructs for analysis of splicing around exon 23 of mutated dystrophin in the mdx mouse mutant, a model for muscular dystrophy. Plans to analyse these constructs in vitro using 2' modified oligonucleotides targeted to splice sites within and adjacent to mouse dystrophin exon 23 were discussed, though no target sites or sequences were given.

2'-O-methyl oligoribonucleotides were subsequently reported to correct dystrophin deficiency in myoblasts from the mdx mouse from this group. An antisense oligonucleotide targeted to the 3' splice site of murine dystrophin intron 22 was reported to cause skipping of the mutant exon as well as several flanking exons and created a novel in-frame dystrophin transcript with a novel internal deletion. This mutated dystrophin was expressed in 1-2% of antisense treated mdx myotubes. Use of other oligonucleotide modifications such as 2'-O-methoxyethyl phosphodiester are described (Dunckley et al. (1998) *Human Mol. Genetics*, 5, 1083-90).

Thus, antisense molecules may provide a tool in the treatment of genetic disorders such as Duchenne Muscular Dystrophy (DMD). However, attempts to induce exon skipping using antisense molecules have had mixed success. Studies on dystrophin exon 19, where successful skipping of that exon from the dystrophin pre-mRNA was achieved using a variety of antisense molecules directed at the flanking splice sites or motifs within the exon involved in exon definition as described by Errington et al. (2003) *J Gen Med* 5, 518-527".

In contrast to the apparent ease of exon 19 skipping, the first report of exon 23 skipping in the mdx mouse by Dunckley et al., (1998) is now considered to be reporting only a naturally occurring revertant transcript or artefact rather than any true antisense activity. In addition to not consistently generating transcripts missing exon 23, Dunckley et al., (1998) did not show any time course of induced exon skipping, or even titration of antisense oligonucleotides, to demonstrate dose dependent effects where the levels of exon skipping corresponded with increasing or decreasing amounts of antisense oligonucleotide. Furthermore, this work could not be replicated by other researchers.

The first example of specific and reproducible exon skipping in the mdx mouse model was reported by Wilton et al., (1999) *Neuromuscular Disorders* 9, 330-338. By directing an antisense molecule to the donor splice site, consistent and efficient exon 23 skipping was induced in the dystrophin mRNA within 6 hours of treatment of the cultured cells. Wilton et al., (1999), also describe targeting the acceptor region of the mouse dystrophin pre-mRNA with longer antisense oligonucleotides and being unable to repeat the published results of Dunckley et al., (1998). No exon skipping, either 23 alone or multiple removal of several flanking exons, could be reproducibly detected using a selection of antisense oligonucleotides directed at the acceptor splice site of intron 22.

While the first antisense oligonucleotide directed at the intron 23 donor splice site induced consistent exon skipping in primary cultured myoblasts, this compound was found to be much less efficient in immortalized cell cultures express-

4

ing higher levels of dystrophin. However, with refined targeting and antisense oligonucleotide design, the efficiency of specific exon removal was increased by almost an order of magnitude (see Mann C J et al., (2002) *J Gen Med* 4, 644-654).

Thus, there remains a need to provide antisense oligonucleotides capable of binding to and modifying the splicing of a target nucleotide sequence. Simply directing the antisense oligonucleotides to motifs presumed to be crucial for splicing is no guarantee of the efficacy of that compound in a therapeutic setting.

SUMMARY OF THE INVENTION

The present invention provides antisense molecule compounds and compositions suitable for binding to RNA motifs involved in the splicing of pre-mRNA that are able to induce specific and efficient exon skipping and a method for their use thereof.

The choice of target selection plays a crucial role in the efficiency of exon skipping and hence its subsequent application of a potential therapy. Simply designing antisense molecules to target regions of pre-mRNA presumed to be involved in splicing is no guarantee of inducing efficient and specific exon skipping. The most obvious or readily defined targets for splicing intervention are the donor and acceptor splice sites although there are less defined or conserved motifs including exonic splicing enhancers, silencing elements and branch points.

The acceptor and donor splice sites have consensus sequences of about 16 and 8 bases respectively (see FIG. 1 for schematic representation of motifs and domains involved in exon recognition, intron removal and the splicing process).

According to a first aspect, the invention provides antisense molecules capable of binding to a selected target to induce exon skipping.

For example, to induce exon skipping in exons 3 to 8, 10 to 16, 19 to 40, 42 to 44, 46, 47, and 50 to 53 in the Dystrophin gene transcript the antisense molecules are preferably selected from the group listed in Table 1A.

In a further example, it is possible to combine two or more antisense oligonucleotides of the present invention together to induce multiple exon skipping in exons 19-20, and 53. This is a similar concept to targeting of a single exon. A combination or "cocktail" of antisense oligonucleotides are directed at adjacent exons to induce efficient exon skipping.

In another example, to induce exon skipping in exons 19-20, 31, 34 and 53 it is possible to improve exon skipping of a single exon by joining together two or more antisense oligonucleotide molecules. This concept is termed by the inventor as a "weasel", an example of a cunningly designed antisense oligonucleotide. A similar concept has been described in Aartsma-Rus A et al., (2004) *Am J Hum Genet.* 74: 83-92).

According to a second aspect, the present invention provides antisense molecules selected and or adapted to aid in the prophylactic or therapeutic treatment of a genetic disorder comprising at least an antisense molecule in a form suitable for delivery to a patient.

According to a third aspect, the invention provides a method for treating a patient suffering from a genetic disease wherein there is a mutation in a gene encoding a particular protein and the affect of the mutation can be abrogated by exon skipping, comprising the steps of: (a) selecting an antisense molecule in accordance with the methods described herein; and (b) administering the molecule to a patient in need of such treatment.

US 8,455,636 B2

5

The invention also addresses the use of purified and isolated antisense oligonucleotides of the invention, for the manufacture of a medicament for treatment of a genetic disease.

The invention further provides a method of treating a condition characterised by Duchenne muscular dystrophy, which method comprises administering to a patient in need of treatment an effective amount of an appropriately designed antisense oligonucleotide of the invention, relevant to the particular genetic lesion in that patient. Further, the invention provides a method for prophylactically treating a patient to prevent or at least minimise Duchene muscular dystrophy, comprising the step of: administering to the patient an effective amount of an antisense oligonucleotide or a pharmaceutical composition comprising one or more of these biological molecules.

The invention also provides kits for treating a genetic disease, which kits comprise at least a antisense oligonucleotide of the present invention, packaged in a suitable container and instructions for its use.

Other aspects and advantages of the invention will become apparent to those skilled in the art from a review of the ensuing description, which proceeds with reference to the following figures.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 Schematic representation of motifs and domains involved in exon recognition, intron removal and the splicing process (SEQ ID NO: 213 and SEQ ID NO: 214).

FIG. 2. Diagrammatic representation of the concept of antisense oligonucleotide induced exon skipping to by-pass disease-causing mutations (not drawn to scale). The hatched box represents an exon carrying a mutation that prevents the translation of the rest of the mRNA into a protein. The solid black bar represents an antisense oligonucleotide that prevents inclusion of that exon in the mature mRNA.

FIG. 3 Gel electrophoresis showing differing efficiencies of two antisense molecules directed at exon 8 acceptor splice site. The preferred compound [H8A(-06+18)] induces strong and consistent exon skipping at a transfection concentration of 20 nanomolar in cultured normal human muscle cells. The less preferred antisense oligonucleotide [H8A(-06+14)] also induces efficient exon skipping, but at much higher concentrations. Other antisense oligonucleotides directed at exon 8 either only induced lower levels of exon skipping or no detectable skipping at all (not shown).

FIG. 4 Gel electrophoresis showing differing efficiencies of two antisense molecules directed at internal domains within exon 7, presumably exon splicing enhancers. The preferred compound [H7A(+45+67)] induces strong and consistent exon skipping at a transfection concentration of 20 nanomolar in cultured human muscle cells. The less preferred antisense oligonucleotide [H7A(+2+26)] induces only low levels of exon skipping at the higher transfection concentrations. Other antisense oligonucleotides directed at exon 7 either only induced lower levels of exon skipping or no detectable skipping at all (not shown).

FIG. 5 Gel electrophoresis showing an example of low efficiency exon 6 skipping using two non-preferred antisense molecules directed at human exon 6 donor splice site. Levels of induced exon 6 skipping are either very low [H6D(+04-21)] or almost undetectable [H6D(+18-04)]. These are examples of non-preferred antisense oligonucleotides to

6

demonstrate that antisense oligonucleotide design plays a crucial role in the efficacy of these compounds.

FIG. 6 Gel electrophoresis showing strong and efficient human exon 6 skipping using an antisense molecules [H6A(+69+91)] directed at an exon 6 internal domain, presumably an exon splicing enhancer. This preferred compound induces consistent exon skipping at a transfection concentration of 20 nanomolar in cultured human muscle cells.

FIG. 7 Gel electrophoresis showing strong human exon 4 skipping using an antisense molecule H4A(+13+32) directed at an exon 6 internal domain, presumably an exon splicing enhancer. This preferred compound induces strong and consistent exon skipping at a transfection concentration of 20 nanomolar in cultured human muscle cells.

FIG. 8 Gel electrophoresis showing (8B) strong human exon 11 skipping using antisense molecule H11A(+75+97) directed at an exon 11 internal domain; and (8B) strong human exon 12 skipping using antisense molecule H12A(+52+75) directed at exon 12 internal domain.

FIG. 9 Gel electrophoresis showing (9A) strong human exon 15 skipping using antisense molecules H15A(+48+71) and H15A(-12+19) directed at an exon 15 internal domain; and (9B) strong human exon 16 skipping using antisense molecules H16A(-12+19) and H16A(-06+25).

FIG. 10 Gel electrophoresis showing human exon 19/20 skipping using antisense molecules H20A(+44+71) and H20A(+149+170) directed at an exon 20 and a "cocktail" of antisense oligonucleotides H19A(+35+65), H20A(+44+71) and H20A(+149+170) directed at exons 19/20.

FIG. 11 Gel electrophoresis showing human exon 19/20 skipping using "weasels" directed at exons 19 and 20.

FIG. 12 Gel electrophoresis showing exon 22 skipping using antisense molecules H22A(+125+106), H22A(+47+69), H22A(+80+101) and H22D(+13-11) directed at exon 22.

FIG. 13 Gel electrophoresis showing exon 31 skipping using antisense molecules H31D(+01-25) and H31D(+03-22); and a "cocktail" of antisense molecules directed at exon 31.

FIG. 14 Gel electrophoresis showing exon 33 skipping using antisense molecules H33A(+30+56) and H33A(+64+88) directed at exon 33.

FIG. 15 Gel electrophoresis showing exon 35 skipping using antisense molecules H35A(+141+161), H35A(+116+135), and H35A(+24+43) and a "cocktail of two antisense molecules, directed at exon 35.

FIG. 16 Gel electrophoresis showing exon 36 skipping using antisense molecules H32A(+49+73) and H36A(+26+50) directed at exon 36.

FIG. 17 Gel electrophoresis showing exon 37 skipping using antisense molecules H37A(+82+105) and H37A(+134+157) directed at exon 37.

FIG. 18 Gel electrophoresis showing exon 38 skipping using antisense molecule H38A(+88+112) directed at exon 38.

FIG. 19 Gel electrophoresis showing exon 40 skipping using antisense molecule H40A(-05+17) directed at exon 40.

FIG. 20 Gel electrophoresis showing exon 42 skipping using antisense molecule H42A(-04+23) directed at exon 42.

FIG. 21 Gel electrophoresis showing exon 46 skipping using antisense molecule H46A(+86+115) directed at exon 46

FIG. 22 Gel electrophoresis showing exon 51, exon 52 and exon 53 skipping using various antisense molecules directed at exons 51, 52 and 53, respectively. A "cocktail" of antisense molecules is also shown directed at exon 53.

US 8,455,636 B2

7

8

TABLE 1A

Description of 2'-O-methyl phosphorothioate antisense oligonucleotides that have been used to date to study induced exon skipping during the processing of the dystrophin pre-mRNA. Since these 2'-O-methyl antisense oligonucleotides are more RNA-like, U represents uracil. With other antisense chemistries such as peptide nucleic acids or morpholinos, these U bases may be shown as "T". Brief Description of the Sequence listings		
SEQ ID	SEQUENCE	NUCLEOTIDE SEQUENCE (5'-3')
1	H8A(-06 + 18)	GAU AGG UGG UAU CAA CAU CUG UAA
2	H8A (-03 + 18)	GAU AGG UGG UAU CAA CAU CUG
3	H8A(-07 + 18)	GAU AGG UGG UAU CAA CAU CUG UAA G
4	H8A(-06 + 14)	GGU GGU AUC AAC AUC UGU AA
5	H8A(-10 + 10)	GUA UCA ACA UCU GUA AGC AC
6	H7A(+45 + 67)	UGC AUG UUC CAG UCG UUG UGU GG
7	H7A(+02 + 26)	CAC UAU UCC AGU CAA AUA GGU CUG G
8	H7D(+15 - 10)	AUU UAC CAA CCU UCA GGA UCG AGU A
9	H7A(-18 + 03)	GGC CUA AAA CAC AUA CAC AUA
10	C6A(-10 + 10)	CAU UUU UGA CCU ACA UGU GG
11	C6A(-14 + 06)	UUU GAC CUA CAU GUG GAA AG
12	C6A(-14 + 12)	UAC AUU UUU GAC CUA CAU GUG GAA AG
13	C6A(-13 + 09)	AUU UUU GAC CUA CAU GGG AAA G
14	CH6A(+69 + 91)	UAC GAG UUG AUU GUC GGA CCC AG
15	C6D(+12 - 13)	GUG GUC UCC UUA CCU AUG ACU GUG G
16	C6D(+06 - 11)	GGU CUC CUU ACC UAU GA
17	H6D(+04 - 21)	UGU CUC AGU AAU CUU CUU ACC UAU
18	H6D(+18 - 04)	UCU UAC CUA UGA CUA UGG AUG AGA
19	H4A(+13 + 32)	GCA UGA ACU CUU GUG GAU CC
20	H4D(+04 - 16)	CCA GGG UAC UAC UUA CAU UA
21	H4D(-24 - 44)	AUC GUG UGU CAC AGC AUC CAG
22	H4A(+11 + 40)	UGU UCA GGG CAU GAA CUC UUG UGG AUC CUU
23	H3A(+30 + 60)	UAG GAG GCG CCU CCC AUC CUG UAG GUC ACU G
24	H3A(+35 + 65)	AGG UCU AGG AGG CGC CUC CCA UCC UGU AGG U
25	H3A(+30 + 54)	GCG CCU CCC AUC CUG UAG GUC ACU G
26	H3D(+46 - 21)	CUU CGA GGA GGU CUA GGA GGC GCC UC
27	H3A(+30 + 50)	CUC CCA UCC UGU AGG UCA CUG
28	H3D(+19 - 03)	UAC CAG UUU UUG CCC UGU CAG G
29	H3A(-06 + 20)	UCA AUA UGC UGC UUC CCA AAC UGA AA
30	H3A(+37 + 61)	CUA GGA GGC GCC UCC CAU CCU GUA G
31	H5A(+20 + 50)	UUA UGA UUU CCA UCU ACG AUG UCA GUA CUU C
32	H5D(+25 - 05)	CUU ACC UGC CAG UGG AGG AUU AUA UUC CAA A

US 8,455,636 B2

9

10

TABLE 1A-continued

Description of 2'-O-methyl phosphorothioate antisense oligonucleotides that have been used to date to study induced exon skipping during the processing of the dystrophin pre-mRNA. Since these 2'-O-methyl antisense oligonucleotides are more RNA-like, U represents uracil. With other antisense chemistries such as peptide nucleic acids or morpholinos, these U bases may be shown as "T".		
Brief Description of the Sequence listings		
SEQ ID	SEQUENCE	NUCLEOTIDE SEQUENCE (5'-3')
33	H5D(+10 - 15)	CAU CAG GAU UCU UAC CUG CCA GUG G
34	H5A(+10 + 34)	CGA UGU CAG UAC UUC CAA UAU UCA C
35	H5D(-04 - 21)	ACC AUU CAU CAG GAU UCU
36	H5D(+16 - 02)	ACC UGC CAG UGG AGG AUU
37	H5A(-07 + 20)	CCA AUA UUC ACU AAA UCA ACC UGU UAA
38	H5D(+18 - 12)	CAG GAU UCU UAC CUG CCA GUG GAG GAU UAU
39	H5A(+05 + 35)	ACG AUG UCA GUA CUU CCA AUA UUC ACU AAA U
40	H5A(+15 + 45)	AUU UCC AUC UAC GAU GUC AGU ACU UCC AAU A
41	H10A(-05 + 16)	CAG GAG CUU CCA AAU GCU GCA
42	H10A(-05 + 24)	CUU GUC UUC AGG AGC UUC CAA AUG CUG CA
43	H10A(+98 + 119)	UCC UCA GCA GAA AGA AGC CAC G
44	H10A(+130 + 149)	UUA GAA AUC UCU CCU UGU GC
45	H10A(-33 - 14)	UAA AUU GGG UGU UAC ACA AU
46	H11D(+26 + 49)	CCC UGA GGC AUU CCC AUC UUG AAU
47	H11D(+11 - 09)	AGG ACU UAC UUG CUU UGU UU
48	H11A(+118 + 140)	CUU GAA UUU AGG AGA UUC AUC UG
49	H11A(+75 + 97)	CAU CUU CUG AUA AUU UUC CUG UU
50	H12A(+52 + 75)	UCU UCU GUU UUU GUU AGC CAG UCA
51	H12A(-10 + 10)	UCU AUG UAA ACU GAA AAU UU
52	H12A(+11 + 30)	UUC UGG AGA UCC AUU AAA AC
53	H13A(+77 + 100)	CAG CAG UUG CGU GAU CUC CAC UAG
54	H13A(+55 + 75)	UUC AUC AAC UAC CAC CAC CAU
55	H13D(+06 - 19)	CUA AGC AAA AUA AUC UGA CCU UAA G
56	H14A(+37 + 64)	CUU GUA AAA GAA CCC AGC GGU CUU CUG U
57	H14A(+14 + 35)	CAU CUA CAG AUG UUU GCC CAU C
58	H14A(+51 + 73)	GAA GGA UGU CUU GUA AAA GAA CC
59	H14D(-02 + 18)	ACC UGU UCU UCA GUA AGA CG
60	H14D(+14 - 10)	CAU GAC ACA CCU GUU CUU CAG UAA
61	H14A(+61 + 80)	CAU UUG AGA AGG AUG UCU UG
62	H14A(-12 + 12)	AUC UCC CAA UAC CUG GAG AAG AGA
63	H15A(-12 + 19)	GCC AUG CAC UAA AAA GGC ACU GCA AGA CAU U
64	H15A(+48 + 71)	UCU UUA AAG CCA GUU GUG UGA AUC
65	H15A(+08 + 28)	UUU CUG AAA GCC AUG CAC UAA

US 8,455,636 B2

11

12

TABLE 1A-continued

Description of 2'-O-methyl phosphorothioate antisense oligonucleotides that have been used to date to study induced exon skipping during the processing of the dystrophin pre-mRNA. Since these 2'-O-methyl antisense oligonucleotides are more RNA-like, U represents uracil. With other antisense chemistries such as peptide nucleic acids or morpholinos, these U bases may be shown as "T".		
Brief Description of the Sequence listings		
SEQ ID	SEQUENCE	NUCLEOTIDE SEQUENCE (5'-3')
66	H15D(+17 - 08)	GUA CAU ACG GCC AGU UUU UGA AGA C
67	H16A(-12 + 19)	CUA GAU CCG CUU UUA AAA CCU GUU AAA ACA A
68	H16A(-06 + 25)	UCU UUU CUA GAU CCG CUU UUA AAA CCU GUU A
69	H16A(-06 + 19)	CUA GAU CCG CUU UUA AAA CCU GUU A
70	H16A(+87 + 109)	CCG UCU UCU GGG UCA CUG ACU UA
71	H16A(-07 + 19)	CUA GAU CCG CUU UUA AAA CCU GUU AA
72	H16A(-07 + 13)	CCG CUU UUA AAA CCU GUU AA
73	H16A(+12 + 37)	UGG AUU GCU UUU UCU UUU CUA GAU CC
74	H16A(+92 + 116)	CAU GCU UCC GUC UUC UGG GUC ACU G
75	H16A(+45 + 67)	G AUC UUG UUU GAG UGA AUA CAG U
76	H16A(+105 + 126)	GUU AUC CAG CCA UGC UUC CGU C
77	H16D(+05 - 20)	UGA UAA UUG GUA UCA CUA ACC UGU G
78	H16D(+12 - 11)	GUA UCA CUA ACC UGU GCU GUA C
79	H19A(+35 + 53)	CUG CUG GCA UCU UGC AGU U
80	H19A(+35 + 65)	GCC UGA GCU GAU CUG CUG GCA UCU UGC AGU U
81	H20A(+44 + 71)	CUG GCA GAA UUC GAU CCA CCG GCU GUU C
82	H20A(+147 + 168)	CAG CAG UAG UUG UCA UCU GCU C
83	H20A(+185 + 203)	UGA UGG GGU GGU GGG UUG G
84	H20A(-08 + 17)	AUC UGC AUU AAC ACC CUC UAG AAA G
85	H20A(+30 + 53)	CCG GCU GUU CAG UUG UUC UGA GGC
86	H20A(-11 + 17)	AUC UGC AUU AAC ACC CUC UAG AAA GAA A
87	H20D(+08 - 20)	GAA GGA GAA GAG AUU CUU ACC UUA CAA A
88	H20A(+44 + 63)	AUU CGA UCC ACC GGC UGU UC
89	H20A(+149 + 168)	CAG CAG UAG UUG UCA UCU GC
90	H21A(-06 + 16)	GCC GGU UGA CUU CAU CCU GUG C
91	H21A(+85 + 106)	CUG CAU CCA GGA ACA UGG GUC C
92	H21A(+85 + 108)	GUC UGC AUC CAG GAA CAU GGG UC
93	H21A(+08 + 31)	GUU GAA GAU CUG AUA GCC GGU UGA
94	H21D(+18 - 07)	UAC UUA CUG UCU GUA GCU CUU UCU
95	H22A(+22 + 45)	CAC UCA UGG UCU CCU GAU AGC GCA
96	H22A(+125 + 146)	CUG CAA UUC CCC GAG UCU CUG C
97	H22A(+47 + 69)	ACU GCU GGA CCC AUG UCC UGA UG
98	H22A(+80 + 101)	CUA AGU UGA GGU AUG GAG AGU
99	H22D(+13 - 11)	UAU UCA CAG ACC UGC AAU UCC CC

US 8,455,636 B2

13

14

TABLE 1A-continued

Description of 2'-O-methyl phosphorothioate antisense oligonucleotides that have been used to date to study induced exon skipping during the processing of the dystrophin pre-mRNA. Since these 2'-O-methyl antisense oligonucleotides are more RNA-like, U represents uracil. With other antisense chemistries such as peptide nucleic acids or morpholinos, these U bases may be shown as "T". Brief Description of the Sequence listings		
SEQ ID	SEQUENCE	NUCLEOTIDE SEQUENCE (5'-3')
100	H23A(+34 + 59)	ACA GUG GUG CUG AGA UAG UAU AGG CC
101	H23A(+18 + 39)	UAG GCC ACU UUG UUG CUC UUG C
102	H23A(+72 + 90)	UUC AGA GGG CGC UUU CUU C
103	H24A(+48 + 70)	GGG CAG GCC AUU CCU CCU UCA GA
104	H24A(-02 + 22)	UCU UCA GGG UUU GUA UGU GAU UCU
105	H25A(+9 + 36)	CUG GGC UGA AUU GUC UGA AUA UCA CUG
106	H25A(+131 + 156)	CUG UUG GCA CAU GUG AUC CCA CUG AG
107	H25D(+16 - 08)	GUC UAU ACC UGU UGG CAC AUG UGA
108	H26A(+132 + 156)	UGC UUU CUG UAA UUC AUC UGG AGU U
109	H26A(-07 + 19)	CCU CCU UUC UGG CAU AGA CCU UCC AC
110	H26A(+68 + 92)	UGU GUC AUC CAU UCG UGC AUC UCU G
111	H27A(+82 + 106)	UUA AGG CCU CUU GUG CUA CAG GUG G
112	H27A(-4 + 19)	GGG CCU CUU CUU UAG CUC UCU GA
113	H27D(+19 - 03)	GAC UUC CAA AGU CUU GCA UUU C
114	H28A(-05 + 19)	GCC AAC AUG CCC AAA CUU CCU AAG
115	H28A(+99 + 124)	CAG AGA UUU CCU CAG CUC CGC CAG GA
116	H28D(+16 - 05)	CUU ACA UCU AGC ACC UCA GAG
117	H29A(+57 + 81)	UCC GCC AUC UGU UAG GGU CUG UGC C
118	H29A(+18 + 42)	AUU UGG GUU AUC CUC UGA AUG UCG C
119	H29D(+17 - 05)	CAU ACC UCU UCA UGU AGU UCC C
120	H30A(+122 + 147)	CAU UUG AGC UGC GUC CAC CUU GUC UG
121	H30A(+25 + 50)	UCC UGG GCA GAC UGG AUG CUC UGU UC
122	H30D(+19 - 04)	UUG CCU GGG CUU CCU GAG GCA UU
123	H31D(+06 - 18)	UUC UGA AAU AAC AUA UAC CUG UGC
124	H31D(+03 - 22)	UAG UUU CUG AAA UAA CAU AUA CCU G
125	H31A(+05 + 25)	GAC UUG UCA AAU CAG AUU GGA
126	H31D(+04 - 20)	GUU UCU GAA AUA ACA UAU ACC UGU
127	H32D(+04 - 16)	CAC CAG AAA UAC AUA CCA CA
128	H32A(+151 + 170)	CAA UGA UUU AGC UGU GAC UG
129	H32A(+10 + 32)	CGA AAC UUC AUG GAG ACA UCU UG
130	H32A(+49 + 73)	CUU GUA GAC GCU GCU CAA AAU UGG C
131	H33D(+09 - 11)	CAU GCA CAC ACC UUU GCU CC
132	H33A(+53 + 76)	UCU GUA CAA UCU GAC GUC CAG UCU
133	H33A(+30 + 56)	GUC UUU AUC ACC AUU UCC ACU UCA GAC

US 8,455,636 B2

15

16

TABLE 1A-continued

Description of 2'-O-methyl phosphorothioate antisense oligonucleotides that have been used to date to study induced exon skipping during the processing of the dystrophin pre-mRNA. Since these 2'-O-methyl antisense oligonucleotides are more RNA-like, U represents uracil. With other antisense chemistries such as peptide nucleic acids or morpholinos, these U bases may be shown as "T".		
Brief Description of the Sequence listings		
SEQ ID	SEQUENCE	NUCLEOTIDE SEQUENCE (5'-3')
134	H33A(+64 + 88)	CCG UCU GCU UUU UCU GUA CAA UCU G
135	H34A(+83 + 104)	UCC AUA UCU GUA GCU GCC AGC C
136	H34A(+143 + 165)	CCA GGC AAC UUC AGA AUC CAA AU
137	H34A(-20 + 10)	UUU CUG UUA CCU GAA AAG AAU UAU AAU GAA
138	H34A(+46 + 70)	CAU UCA UUU CCU UUC GCA UCU UAC G
139	H34A(+95 + 120)	UGA UCU CUU UGU CAA UUC CAU AUC UG
140	H34D(+10 - 20)	UUC AGU GAU AUA GGU UUU ACC UUU CCC CAG
141	H34A(+72 + 96)	CUG UAG CUG CCA GCC AUU CUG UCA AG
142	H35A(+141 + 161)	UCU UCU GCU CGG GAG GUG ACA
143	H35A(+116 + 135)	CCA GUU ACU AUU CAG AAG AC
144	H35A(+24 + 43)	UCU UCA GGU GCA CCU UCU GU
145	H36A(+26 + 50)	UGU GAU GUG GUC CAC AUU CUG GUC A
146	H36A(-02 + 18)	CCA UGU GUU UCU GGU AUU CC
147	H37A(+26 + 50)	CGU GUA GAG UCC ACC UUU GGG CGU A
148	H37A(+82 + 105)	UAC UAA UUU CCU GCA GUG GUC ACC
149	H37A(+134 + 157)	UUC UGU GUG AAA UGG CUG CAA AUC
150	H38A(-01 + 19)	CCU UCA AAG GAA UGG AGG CC
151	H38A(+59 + 83)	UGC UGA AUU UCA GCC UCC AGU GGU U
152	H38A(+88 + 112)	UGA AGU CUU CCU CUU UCA GAU UCA C
153	H39A(+62 + 85)	CUG GCU UUC UCU CAU CUG UGA UUC
154	H39A(+39 + 58)	GUU GUA AGU UGU CUC CUC UU
155	H39A(+102 + 121)	UUG UCU GUA ACA GCU GCU GU
156	H39D(+10 - 10)	GCU CUA AUA CCU UGA GAG CA
157	H40A(-05 + 17)	CUU UGA GAC CUC AAA UCC UGU U
158	H40A(+129 + 153)	CUU UAU UUU CCU UUC AUC UCU GGG C
159	H42A(-04 + 23)	AUC GUU UCU UCA CGG ACA GUG UGC UGG
160	H42A(+86 + 109)	GGG CUU GUG AGA CAU GAG UGA UUU
161	H42D(+19 - 02)	A CCU UCA GAG GAC UCC UCU UGC
162	H43D(+10 - 15)	UAU GUG UUA CCU ACC CUU GUC GGU C
163	H43A(+101 + 120)	GGA GAG AGC UUC CUG UAG CU
164	H43A(+78 + 100)	UCA CCC UUU CCA CAG GCG UUG CA
165	H44A(+85 + 104)	UUU GUG UCU UUC UGA GAA AC
166	H44D(+10 - 10)	AAA GAC UUA CCU UAA GAU AC
167	H44A(-06 + 14)	AUC UGU CAA AUC GCC UGC AG

US 8,455,636 B2

17

18

TABLE 1A-continued

Description of 2'-O-methyl phosphorothioate antisense oligonucleotides that have been used to date to study induced exon skipping during the processing of the dystrophin pre-mRNA. Since these 2'-O-methyl antisense oligonucleotides are more RNA-like, U represents uracil. With other antisense chemistries such as peptide nucleic acids or morpholinos, these U bases may be shown as "T".		
Brief Description of the Sequence listings		
SEQ ID	SEQUENCE	NUCLEOTIDE SEQUENCE (5'-3')
168	H46D(+16 - 04)	UUA CCU UGA CUU GCU CAA GC
169	H46A(+90 + 109)	UCC AGG UUC AAG UGG GAU AC
170	H47A(+76 + 100)	GCU CUU CUG GGC UUA UGG GAG CAC U
171	H47D(+25 - 02)	ACC UUU AUC CAC UGG AGA UUU GUC UGC
172	H47A(-9 + 12)	UUC CAC CAG UAA CUG AAA CAG
173	H50A(+02 + 30)	CCA CUC AGA GCU CAG AUC UUC UAA CUU CC
174	H50A(+07 + 33)	CUU CCA CUC AGA GCU CAG AUC UUC UAA
175	H50D(+07 - 18)	GGG AUC CAG UAU ACU UAC AGG CUC C
176	H51A(-01 + 25)	ACC AGA GUA ACA GUC UGA GUA GGA GC
177	H51D(+16 - 07)	CUC AUA CCU UCU GCU UGA UGA UC
178	H51A(+111 + 134)	UUC UGU CCA AGC CCG GUU GAA AUC
179	H51A(+61 + 90)	ACA UCA AGG AAG AUG GCA UUU CUA GUU UGG
180	H51A(+66 + 90)	ACA UCA AGG AAG AUG GCA UUU CUA G
181	H51A(+66 + 95)	CUC CAA CAU CAA GGA AGA UGG CAU UUC UAG
182	H51D(+08 - 17)	AUC AUU UUU UCU CAU ACC UUC UGC U
183	H51A/D(+08 - 17 & (-15+))	AUC AUU UUU UCU CAU ACC UUC UGC UAG GAG CUA AAA
184	H51A(+175 + 195)	CAC CCA CCA UCA CCC UCU GUG
185	H51A(+199 + 220)	AUC AUC UCG UUG AUA UCC UCA A
186	H52A(-07 + 14)	UCC UGC AUU GUU GCC UGU AAG
187	H52A(+12 + 41)	UCC AAC UGG GGA CGC CUC UGU UCC AAA UCC
188	H52A(+17 + 37)	ACU GGG GAC GCC UCU GUU CCA
189	H52A(+93 + 112)	CCG UAA UGA UUG UUC UAG CC
190	H52D(+05 - 15)	UGU UAA AAA ACU UAC UUC GA
191	H53A(+45 + 69)	CAU UCA ACU GUU GCC UCC GGU UCU G
192	H53A(+39 + 62)	CUG UUG CCU CCG GUU CUG AAG GUG
193	H53A(+39 + 69)	CAU UCA ACU GUU GCC UCC GGU UCU GAA GGU G
194	H53D(+14 - 07)	UAC UAA CCU UGG UUU CUG UGA
195	H53A(+23 + 47)	CUG AAG GUG UUC UUG UAC UUC AUC C
196	H53A(+150 + 176)	UGU AUA GGG ACC CUC CUU CCA UGA CUC
197	H53D(+20 - 05)	CUA ACC UUG GUU UCU GUG AUU UUC U
198	H53D(+09 - 18)	GGU AUC UUU GAU ACU AAC CUU GGU UUC
199	H53A(-12 + 10)	AUU CUU UCA ACU AGA AUA AAA G

US 8,455,636 B2

19

TABLE 1A-continued

Description of 2'-O-methyl phosphorothioate antisense oligonucleotides that have been used to date to study induced exon skipping during the processing of the dystrophin pre-mRNA. Since these 2'-O-methyl antisense oligonucleotides are more RNA-like, U represents uracil. With other antisense chemistries such as peptide nucleic acids or morpholinos, these U bases may be shown as "T". Brief Description of the Sequence listings		
SEQ ID	SEQUENCE	NUCLEOTIDE SEQUENCE (5'-3')
200	H53A(-07 + 18)	GAU UCU GAA UUC UUU CAA CUA GAA U
201	H53A(+07 + 26)	AUC CCA CUG AUU CUG AAU UC
202	H53A(+124 + 145)	UUG GCU CUG GCC UGU CCU AAG A
203	H46A(+86 + 115)	CUC UUU UCC AGG UUC AAG UGG GAU ACU AGC
204	H46A(+107 + 137)	CAA GCU UUU CUU UUA GUU GCU GCU CUU UUC C
205	H46A(-10 + 20)	UAU UCU UUU GUU CUU CUA GCC UGG AGA AAG
206	H46A(+50 + 77)	CUG CUU CCU CCA ACC AUA AAA CAA AUU C
207	H45A(-06 + 20)	CCA AUG CCA UCC UGG AGU UCC UGU AA
208	H45A(+91 + 110)	UCC UGU AGA AUA CUG GCA UC
209	H45A(+125 + 151)	UGC AGA CCU CCU GCC ACC GCA GAU UCA
210	H45D(+16 - 04)	CUA CCU CUU UUU UCU GUC UG
211	H45A(+71 + 90)	UGU UUU UGA GGA UUG CUG AA

20

TABLE 1B

Description of a cocktail of 2'-O-methyl phosphorothioate antisense oligonucleotides that have been used to date to study induced exon skipping during the processing of the dystrophin pre-mRNA.		
SEQ ID	SEQUENCE	NUCLEOTIDE SEQUENCE (5'-3')
81	H20A(+44 + 71)	CUG GCA GAA UUC GAU CCA CCG GCU GUU C
82	H20A(+147 + 168)	CAG CAG UAG UUG UCA UCU GCU C
80	H19A(+35 + 65)	GCC UGA GCU GAU CUG CUG GCA UCU UGC AGU U
81	H20A(+44 + 71)	CUG GCA GAA UUC GAU CCA CCG GCU GUU C

35

TABLE 1B-continued

Description of a cocktail of 2'-O-methyl phosphorothioate antisense oligonucleotides that have been used to date to study induced exon skipping during the processing of the dystrophin pre-mRNA.		
SEQ ID	SEQUENCE	NUCLEOTIDE SEQUENCE (5'-3')
82	H20A(+147 + 168)	CAG CAG UAG UUG UCA UCU GCU C
194	H53D(+14 - 07)	UAC UAA CCU UGG UUU CUG UGA
195	H53A(+23 + 47)	CUG AAG GUG UUC UUG UAC UUC AUC C
196	H53A(+150 + 176)	UGU AUA GGG ACC CUC CUU CCA UGA CUC

40

45

TABLE 1C

Description of a "weasel" of 2'-O-methyl phosphorothioate antisense oligonucleotides that have been used to date to study induced exon skipping during the processing of the dystrophin pre-mRNA.	
SEQ ID	SEQUENCE
SEQ ID	SEQUENCE
81 H20A(+44 + 71) -	82 H20A(+147 + 168)
CUG GCA GAA UUC GAU CCA CCG GCU GUU C-	CAG CAG UAG UUG UCA UCU GCU C
80 H19A(+35 + 65) -	88 H20A(+44 + 63) -
GCC UGA GCU GAU CUG CUG GCA UCU UGC AGU U	-AUU CGA UCC ACC GGC UGU UC-
79 H20A(+149 + 168)	
CUG CUG GCA UCU UGC AGU U	

US 8,455,636 B2

21

22

TABLE 1C-continued

Description of a "weasel" of 2'-O-methyl phosphorothioate antisense oligonucleotides that have been used to date to study induced exon skipping during the processing of the dystrophin pre-mRNA.	
SEQ ID SEQUENCE	NUCLEOTIDE SEQUENCE (5'-3')
80H19A(+35 + 65) -	GCC UGA GCU GAU CUG CUG GCA UCU UGC AGU U
88H20A(+44 + 63)	-AUU CGA UCC ACC GGC UGU UC-
80H19A(+35 + 65) -	GCC UGA GCU GAU CUG CUG GCA UCU UGC AGU U
79H20A(+149 + 168)	-CUG CUG GCA UCU UGC AGU U
138H34A(+46 + 70) -	CAU UCA UUU CCU UUC GCA UCU UAC G-
139H34A(+94 + 120)	UGA UCU CUU UGU CAA UUC CAU AUC UG
124H31D(+03 - 22) -	UAG UUU CUG AAA UAA CAU AUA CCU G-
144UU- H35A(+24 + 43)	UU- UCU UCA GGU GCA CCU UCU GU
195H53A(+23 + 47) -	CUG AAG GUG UUC UUG UAC UUC AUC C-
AA-	
196H53A(+150 + 176) -	UGU AUA GGG ACC CUC CUU CCA
AA-	UGA CUC-AA-
194H53D(+14 - 07)	UAC UAA CCU UGG UUU CUG UGA
212Aimed at exons 19/20/20	CAG CAG UAG UUG UCA UCU GCU CAA CUG GCA GAA UUC GAU CCA CCG GCU GUU CAA GCC UGA GCU GAU CUG CUC GCA UCU UGC AGU

DETAILED DESCRIPTION OF THE INVENTION

General

Those skilled in the art will appreciate that the invention described herein is susceptible to variations and modifications other than those specifically described. It is to be understood that the invention includes all such variation and modifications. The invention also includes all of the steps, features, compositions and compounds referred to or indicated in the specification, individually or collectively and any and all combinations or any two or more of the steps or features.

The present invention is not to be limited in scope by the specific embodiments described herein, which are intended for the purpose of exemplification only. Functionally equivalent products, compositions and methods are clearly within the scope of the invention as described herein.

Sequence identity numbers (SEQ ID NO:) containing nucleotide and amino acid sequence information included in this specification are collected at the end of the description and have been prepared using the programme PatentIn Version 3.0. Each nucleotide or amino acid sequence is identified in the sequence listing by the numeric indicator <210> followed by the sequence identifier (e.g. <210>1, <210>2, etc.). The length, type of sequence and source organism for each nucleotide or amino acid sequence are indicated by information provided in the numeric indicator fields <211>, <212> and <213>, respectively. Nucleotide and amino acid sequences referred to in the specification are defined by the information provided in numeric indicator field <400> followed by the sequence identifier (e.g. <400>1, <400>2, etc.).

An antisense molecules nomenclature system was proposed and published to distinguish between the different antisense molecules (see Mann et al., (2002) *J Gen Med* 4, 644-654). This nomenclature became especially relevant when

testing several slightly different antisense molecules, all directed at the same target region, as shown below:

H#A/D(x;y).

The first letter designates the species (e.g. H: human, M: murine, C: canine)

"#" designates target dystrophin exon number.

"A/D" indicates acceptor or donor splice site at the beginning and end of the exon, respectively.

(x;y) represents the annealing coordinates where "-" or "+" indicate intronic or exonic sequences respectively. As an example, A(-6+18) would indicate the last 6 bases of the intron preceding the target exon and the first 18 bases of the target exon. The closest splice site would be the acceptor so these coordinates would be preceded with an "A". Describing annealing coordinates at the donor splice site could be D(+2-18) where the last 2 exonic bases and the first 18 intronic bases correspond to the annealing site of the antisense molecule. Entirely exonic annealing coordinates that would be represented by A(+65+85), that is the site between the 65th and 85th nucleotide from the start of that exon.

The entire disclosures of all publications (including patents, patent applications, journal articles, laboratory manuals, books, or other documents) cited herein are hereby incorporated by reference. No admission is made that any of the references constitute prior art or are part of the common general knowledge of those working in the field to which this invention relates.

As used necessarily herein the term "derived" and "derived from" shall be taken to indicate that a specific integer may be obtained from a particular source albeit not directly from that source.

Throughout this specification, unless the context requires otherwise, the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the

US 8,455,636 B2

23

inclusion of a stated integer or group of integers but not the exclusion of any other integer or group of integers.

Other definitions for selected terms used herein may be found within the detailed description of the invention and apply throughout. Unless otherwise defined, all other scientific and technical terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which the invention belongs.

DESCRIPTION OF THE PREFERRED EMBODIMENT

When antisense molecule(s) are targeted to nucleotide sequences involved in splicing in exons within pre-mRNA sequences, normal splicing of the exon may be inhibited causing the splicing machinery to by-pass the entire mutated exon from the mature mRNA. The concept of antisense oligonucleotide induced exon skipping is shown in FIG. 2. In many genes, deletion of an entire exon would lead to the production of a non-functional protein through the loss of important functional domains or the disruption of the reading frame. In some proteins, however, it is possible to shorten the protein by deleting one or more exons, without disrupting the reading frame, from within the protein without seriously altering the biological activity of the protein. Typically, such proteins have a structural role and/or possess functional domains at their ends. The present invention describes antisense molecules capable of binding to specified dystrophin pre-mRNA targets and re-directing processing of that gene. Antisense Molecules

According to a first aspect of the invention, there is provided antisense molecules capable of binding to a selected target to induce exon skipping. To induce exon skipping in exons of the Dystrophin gene transcript, the antisense molecules are preferably selected from the group of compounds shown in Table 1A. There is also provided a combination or "cocktail" of two or more antisense oligonucleotides capable of binding to a selected target to induce exon skipping. To induce exon skipping in exons of the Dystrophin gene transcript, the antisense molecules in a "cocktail" are preferably selected from the group of compounds shown in Table 1B. Alternatively, exon skipping may be induced by antisense oligonucleotides joined together "weasels" preferably selected from the group of compounds shown in Table 1C.

Designing antisense molecules to completely mask consensus splice sites may not necessarily generate any skipping of the targeted exon. Furthermore, the inventors have discovered that size or length of the antisense oligonucleotide itself is not always a primary factor when designing antisense molecules. With some targets such as exon 19, antisense oligonucleotides as short as 12 bases were able to induce exon skipping, albeit not as efficiently as longer (20-31 bases) oligonucleotides. In some other targets, such as murine dystrophin exon 23, antisense oligonucleotides only 17 residues long were able to induce more efficient skipping than another overlapping compound of 25 nucleotides.

The inventors have also discovered that there does not appear to be any standard motif that can be blocked or masked by antisense molecules to redirect splicing. In some exons, such as mouse dystrophin exon 23, the donor splice site was the most amenable to target to re-direct skipping of that exon. It should be noted that designing and testing a series of exon 23 specific antisense molecules to anneal to overlapping regions of the donor splice site showed considerable variation in the efficacy of induced exon skipping. As reported in Mann et al., (2002) there was a significant variation in the efficiency of bypassing the nonsense mutation depending upon anti-

24

sense oligonucleotide annealing ("Improved antisense oligonucleotide induced exon skipping in the mdx mouse model of muscular dystrophy". *J Gen Med* 4: 644-654). Targeting the acceptor site of exon 23 or several internal domains was not found to induce any consistent exon 23 skipping.

In other exons targeted for removal, masking the donor splice site did not induce any exon skipping. However, by directing antisense molecules to the acceptor splice site (human exon 8 as discussed below), strong and sustained exon skipping was induced. It should be noted that removal of human exon 8 was tightly linked with the co-removal of exon 9. There is no strong sequence homology between the exon 8 antisense oligonucleotides and corresponding regions of exon 9 so it does not appear to be a matter of cross reaction. Rather the splicing of these two exons is inextricably linked. This is not an isolated instance as the same effect is observed in canine cells where targeting exon 8 for removal also resulted in the skipping of exon 9. Targeting exon 23 for removal in the mouse dystrophin pre-mRNA also results in the frequent removal of exon 22 as well. This effect occurs in a dose dependent manner and also indicates close coordinated processing of 2 adjacent exons.

In other targeted exons, antisense molecules directed at the donor or acceptor splice sites did not induce exon skipping while annealing antisense molecules to intra-exonic regions (i.e. exon splicing enhancers within human dystrophin exon 6) was most efficient at inducing exon skipping. Some exons, both mouse and human exon 19 for example, are readily skipped by targeting antisense molecules to a variety of motifs. That is, targeted exon skipping is induced after using antisense oligonucleotides to mask donor and acceptor splice sites or exon splicing enhancers.

To identify and select antisense oligonucleotides suitable for use in the modulation of exon skipping, a nucleic acid sequence whose function is to be modulated must first be identified. This may be, for example, a gene (or mRNA transcribed from the gene) whose expression is associated with a particular disorder or disease state, or a nucleic acid molecule from an infectious agent. Within the context of the present invention, preferred target site(s) are those involved in mRNA splicing (i.e. splice donor sites, splice acceptor sites, or exonic splicing enhancer elements). Splicing branch points and exon recognition sequences or splice enhancers are also potential target sites for modulation of mRNA splicing.

Preferably, the present invention aims to provide antisense molecules capable of binding to a selected target in the dystrophin pre-mRNA to induce efficient and consistent exon skipping. Duchenne muscular dystrophy arises from mutations that preclude the synthesis of a functional dystrophin gene product. These Duchenne muscular dystrophy gene defects are typically nonsense mutations or genomic rearrangements such as deletions, duplications or micro-deletions or insertions that disrupt the reading frame. As the human dystrophin gene is a large and complex gene with the 79 exons being spliced together to generate a mature mRNA with an open reading frame of approximately 11,000 bases, there are many positions where these mutations can occur. Consequently, a comprehensive antisense oligonucleotide based therapy to address many of the different disease-causing mutations in the dystrophin gene will require that many exons can be targeted for removal during the splicing process.

Within the context of the present invention, preferred target site(s) are those involved in mRNA splicing (i.e. splice donor sites, splice acceptor sites or exonic splicing enhancer elements). Splicing branch points and exon recognition sequences or splice enhancers are also potential target sites for modulation of mRNA splicing.

US 8,455,636 B2

25

The oligonucleotide and the DNA or RNA are complementary to each other when a sufficient number of corresponding positions in each molecule are occupied by nucleotides which can hydrogen bond with each other. Thus, "specifically hybridisable" and "complementary" are terms which are used to indicate a sufficient degree of complementarity or precise pairing such that stable and specific binding occurs between the oligonucleotide and the DNA or RNA target. It is understood in the art that the sequence of an antisense molecule need not be 100% complementary to that of its target sequence to be specifically hybridisable. An antisense molecule is specifically hybridisable when binding of the compound to the target DNA or RNA molecule interferes with the normal function of the target DNA or RNA to cause a loss of utility, and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target sequences under conditions in which specific binding is desired, i.e., under physiological conditions in the case of in vivo assays or therapeutic treatment, and in the case of in vitro assays, under conditions in which the assays are performed.

While the above method may be used to select antisense molecules capable of deleting any exon from within a protein that is capable of being shortened without affecting its biological function, the exon deletion should not lead to a reading frame shift in the shortened transcribed mRNA. Thus, if in a linear sequence of three exons the end of the first exon encodes two of three nucleotides in a codon and the next exon is deleted then the third exon in the linear sequence must start with a single nucleotide that is capable of completing the nucleotide triplet for a codon. If the third exon does not commence with a single nucleotide there will be a reading frame shift that would lead to the generation of truncated or a non-functional protein.

It will be appreciated that the codon arrangements at the end of exons in structural proteins may not always break at the end of a codon, consequently there may be a need to delete more than one exon from the pre-mRNA to ensure in-frame reading of the mRNA. In such circumstances, a plurality of antisense oligonucleotides may need to be selected by the method of the invention wherein each is directed to a different region responsible for inducing splicing in the exons that are to be deleted.

The length of an antisense molecule may vary so long as it is capable of binding selectively to the intended location within the pre-mRNA molecule. The length of such sequences can be determined in accordance with selection procedures described herein. Generally, the antisense molecule will be from about 10 nucleotides in length up to about 50 nucleotides in length. It will be appreciated however that any length of nucleotides within this range may be used in the method. Preferably, the length of the antisense molecule is between 17 to 30 nucleotides in length.

In order to determine which exons can be connected in a dystrophin gene, reference should be made to an exon boundary map. Connection of one exon with another is based on the exons possessing the same number at the 3' border as is present at the 5' border of the exon to which it is being connected. Therefore, if exon 7 were deleted, exon 6 must connect to either exons 12 or 18 to maintain the reading frame. Thus, antisense oligonucleotides would need to be selected which redirected splicing for exons 7 to 11 in the first instance or exons 7 to 17 in the second instance. Another and somewhat simpler approach to restore the reading frame around an exon 7 deletion would be to remove the two flanking exons. Induction of exons 6 and 8 skipping should result in an in-frame transcript with the splicing of exons 5 to 9. In practise however, targeting exon 8 for removal from the pre-

26

mRNA results in the co-removal of exon 9 so the resultant transcript would have exon 5 joined to exon 10. The inclusion or exclusion of exon 9 does not alter the reading frame. Once the antisense molecules to be tested have been identified, they are prepared according to standard techniques known in the art. The most common method for producing antisense molecules is the methylation of the 2' hydroxyribose position and the incorporation of a phosphorothioate backbone produces molecules that superficially resemble RNA but that are much more resistant to nuclease degradation.

To avoid degradation of pre-mRNA during duplex formation with the antisense molecules, the antisense molecules used in the method may be adapted to minimise or prevent cleavage by endogenous RNase H. This property is highly preferred as the treatment of the RNA with the unmethylated oligonucleotides either intracellularly or in crude extracts that contain RNase H leads to degradation of the pre-mRNA: antisense oligonucleotide duplexes. Any form of modified antisense molecules that is capable of by-passing or not inducing such degradation may be used in the present method. An example of antisense molecules which when duplexed with RNA are not cleaved by cellular RNase H is 2'-O-methyl derivatives. 2'-O-methyl-oligoribonucleotides are very stable in a cellular environment and in animal tissues, and their duplexes with RNA have higher T_m values than their ribo- or deoxyribo-counterparts.

Antisense molecules that do not activate RNase H can be made in accordance with known techniques (see, e.g., U.S. Pat. No. 5,149,797). Such antisense molecules, which may be deoxyribonucleotide or ribonucleotide sequences, simply contain any structural modification which sterically hinders or prevents binding of RNase H to a duplex molecule containing the oligonucleotide as one member thereof, which structural modification does not substantially hinder or disrupt duplex formation. Because the portions of the oligonucleotide involved in duplex formation are substantially different from those portions involved in RNase H binding thereto, numerous antisense molecules that do not activate RNase H are available. For example, such antisense molecules may be oligonucleotides wherein at least one, or all, of the inter-nucleotide bridging phosphate residues are modified phosphates, such as methyl phosphonates, methyl phosphorothioates, phosphoromorpholidates, phosphoropiperazidates and phosphoramidates. For example, every other one of the internucleotide bridging phosphate residues may be modified as described. In another non-limiting example, such antisense molecules are molecules wherein at least one, or all, of the nucleotides contain a 2' lower alkyl moiety (e.g., C₁-C₄, linear or branched, saturated or unsaturated alkyl, such as methyl, ethyl, ethenyl, propyl, 1-propenyl, 2-propenyl, and isopropyl). For example, every other one of the nucleotides may be modified as described.

While antisense oligonucleotides are a preferred form of the antisense molecules, the present invention comprehends other oligomeric antisense molecules, including but not limited to oligonucleotide mimetics such as are described below.

Specific examples of preferred antisense compounds useful in this invention include oligonucleotides containing modified backbones or non-natural inter-nucleoside linkages. As defined in this specification, oligonucleotides having modified backbones include those that retain a phosphorus atom in the backbone and those that do not have a phosphorus atom in the backbone. For the purposes of this specification, and as sometimes referenced in the art, modified oligonucleotides that do not have a phosphorus atom in their internucleoside backbone can also be considered to be oligonucleosides.

In other preferred oligonucleotide mimetics, both the sugar and the inter-nucleoside linkage, i.e., the backbone, of the nucleotide units are replaced with novel groups. The base units are maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound, an oligonucleotide mimetic that has been shown to have excellent hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar-backbone of an oligonucleotide is replaced with an amide containing backbone, in particular an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone.

Modified oligonucleotides may also contain one or more substituted sugar moieties. Oligonucleotides may also include nucleobase (often referred to in the art simply as "base") modifications or substitutions. Certain nucleobases are particularly useful for increasing the binding affinity of the oligomeric compounds of the invention. These include 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, including 2-aminopropyladenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2° C. and are presently preferred base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar modifications.

Another modification of the oligonucleotides of the invention involves chemically linking to the oligonucleotide one or more moieties or conjugates that enhance the activity, cellular distribution or cellular uptake of the oligonucleotide. Such moieties include but are not limited to lipid moieties such as a cholesterol moiety, cholic acid, a thioether, e.g., hexyl-5-tritylthiol, a thiocholesterol, an aliphatic chain, e.g., dodecanol or undecyl residues, a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate, a polyamine or a polyethylene glycol chain, or adamantane acetic acid, a palmityl moiety, or an octadecylamine or hexylamino-carbonyl-oxysterol moiety.

It is not necessary for all positions in a given compound to be uniformly modified, and in fact more than one of the aforementioned modifications may be incorporated in a single compound or even at a single nucleoside within an oligonucleotide. The present invention also includes antisense compounds that are chimeric compounds. "Chimeric" antisense compounds or "chimeras," in the context of this invention, are antisense molecules, particularly oligonucleotides, which contain two or more chemically distinct regions, each made up of at least one monomer unit, i.e., a nucleotide in the case of an oligonucleotide compound. These oligonucleotides typically contain at least one region wherein the oligonucleotide is modified so as to confer upon the increased resistance to nuclease degradation, increased cellular uptake, and an additional region for increased binding affinity for the target nucleic acid.

Methods of Manufacturing Antisense Molecules

The antisense molecules used in accordance with this invention may be conveniently and routinely made through the well-known technique of solid phase synthesis. Equipment for such synthesis is sold by several vendors including, for example, Applied Biosystems (Foster City, Calif.). One method for synthesizing oligonucleotides on a modified solid support is described in U.S. Pat. No. 4,458,066.

Any other means for such synthesis known in the art may additionally or alternatively be employed. It is well known to use similar techniques to prepare oligonucleotides such as the phosphorothioates and alkylated derivatives. In one such

automated embodiment, diethyl-phosphoramidites are used as starting materials and may be synthesized as described by Beaucage, et al., (1981) *Tetrahedron Letters*, 22:1859-1862.

The antisense molecules of the invention are synthesized *in vitro* and do not include antisense compositions of biological origin, or genetic vector constructs designed to direct the *in vivo* synthesis of antisense molecules. The molecules of the invention may also be mixed, encapsulated, conjugated or otherwise associated with other molecules, molecule structures or mixtures of compounds, as for example, liposomes, receptor targeted molecules, oral, rectal, topical or other formulations, for assisting in uptake, distribution and/or absorption.

Therapeutic Agents

The present invention also can be used as a prophylactic or therapeutic, which may be utilized for the purpose of treatment of a genetic disease.

Accordingly, in one embodiment the present invention provides antisense molecules that bind to a selected target in the dystrophin pre-mRNA to induce efficient and consistent exon skipping described herein in a therapeutically effective amount admixed with a pharmaceutically acceptable carrier, diluent, or excipient.

The phrase "pharmaceutically acceptable" refers to molecular entities and compositions that are physiologically tolerable and do not typically produce an allergic or similarly untoward reaction, such as gastric upset and the like, when administered to a patient. The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the compound is administered. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water or saline solutions and aqueous dextrose and glycerol solutions are preferably employed as carriers, particularly for injectable solutions. Suitable pharmaceutical carriers are described in Martin, *Remington's Pharmaceutical Sciences*, 18th Ed., Mack Publishing Co., Easton, Pa., (1990).

In a more specific form of the invention there are provided pharmaceutical compositions comprising therapeutically effective amounts of an antisense molecule together with pharmaceutically acceptable diluents, preservatives, solubilizers, emulsifiers, adjuvants and/or carriers. Such compositions include diluents of various buffer content (e.g., Tris-HCl, acetate, phosphate), pH and ionic strength and additives such as detergents and solubilizing agents (e.g., Tween 80, Polysorbate 80), anti-oxidants (e.g., ascorbic acid, sodium metabisulfite), preservatives (e.g., Thimersol, benzyl alcohol) and bulking substances (e.g., lactose, mannitol). The material may be incorporated into particulate preparations of polymeric compounds such as polylactic acid, polyglycolic acid, etc. or into liposomes. Hylauronic acid may also be used. Such compositions may influence the physical state, stability, rate of *in vivo* release, and rate of *in vivo* clearance of the present proteins and derivatives. See, e.g., Martin, *Remington's Pharmaceutical Sciences*, 18th Ed. (1990), Mack Publishing Co., Easton, Pa. 18042) pages 1435-1712 that are herein incorporated by reference. The compositions may be prepared in liquid form, or may be in dried powder, such as lyophilized form.

It will be appreciated that pharmaceutical compositions provided according to the present invention may be administered by any means known in the art. Preferably, the pharmaceutical compositions for administration are administered by injection, orally, or by the pulmonary, or nasal route. The antisense molecules are more preferably delivered by intra-

US 8,455,636 B2

29

venous, intra-arterial, intraperitoneal, intramuscular, or subcutaneous routes of administration.

Antisense Molecule Based Therapy

Also addressed by the present invention is the use of antisense molecules of the present invention, for manufacture of a medicament for modulation of a genetic disease.

The delivery of a therapeutically useful amount of antisense molecules may be achieved by methods previously published. For example, intracellular delivery of the antisense molecule may be via a composition comprising an admixture of the antisense molecule and an effective amount of a block copolymer. An example of this method is described in US patent application US 20040248833.

Other methods of delivery of antisense molecules to the nucleus are described in Mann C J et al., (2001) [*"Antisense-induced exon skipping and the synthesis of dystrophin in the mdx mouse"*]. Proc., Natl. Acad. Science, 98(1) 42-47] and in Gebiski et al., (2003). Human Molecular Genetics, 12(15): 1801-1811.

A method for introducing a nucleic acid molecule into a cell by way of an expression vector either as naked DNA or complexed to lipid carriers, is described in U.S. Pat. No. 6,806,084.

It may be desirable to deliver the antisense molecule in a colloidal dispersion system. Colloidal dispersion systems include macromolecule complexes, nanocapsules, microspheres, beads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes or liposome formulations.

Liposomes are artificial membrane vesicles which are useful as delivery vehicles in vitro and in vivo. These formulations may have net cationic, anionic or neutral charge characteristics and are useful characteristics with in vitro, in vivo and ex vivo delivery methods. It has been shown that large unilamellar vesicles (LUV), which range in size from 0.2-4.0.µm can encapsulate a substantial percentage of an aqueous buffer containing large macromolecules. RNA, and DNA can be encapsulated within the aqueous interior and be delivered to cells in a biologically active form (Fraley, et al., Trends Biochem. Sci., 6:77, 1981).

In order for a liposome to be an efficient gene transfer vehicle, the following characteristics should be present: (1) encapsulation of the antisense molecule of interest at high efficiency while not compromising their biological activity; (2) preferential and substantial binding to a target cell in comparison to non-target cells; (3) delivery of the aqueous contents of the vesicle to the target cell cytoplasm at high efficiency; and (4) accurate and effective expression of genetic information (Mannino, et al., Biotechniques, 6:682, 1988).

The composition of the liposome is usually a combination of phospholipids, particularly high-phase-transition-temperature phospholipids, usually in combination with steroids, especially cholesterol. Other phospholipids or other lipids may also be used. The physical characteristics of liposomes depend on pH, ionic strength, and the presence of divalent cations.

Alternatively, the antisense construct may be combined with other pharmaceutically acceptable carriers or diluents to produce a pharmaceutical composition. Suitable carriers and diluents include isotonic saline solutions, for example phosphate-buffered saline. The composition may be formulated for parenteral, intramuscular, intravenous, subcutaneous, intraocular, oral or transdermal administration.

The routes of administration described are intended only as a guide since a skilled practitioner will be able to determine readily the optimum route of administration and any dosage

30

for any particular animal and condition. Multiple approaches for introducing functional new genetic material into cells, both in vitro and in vivo have been attempted (Friedmann (1989) Science, 244:1275-1280). These approaches include integration of the gene to be expressed into modified retroviruses (Friedmann (1989) supra; Rosenberg (1991) Cancer Research 51(18), suppl.: 5074S-5079S); integration into non-retrovirus vectors (Rosenfeld, et al. (1992) Cell, 68:143-155; Rosenfeld, et al. (1991) Science, 252:431-434); or delivery of a transgene linked to a heterologous promoter-enhancer element via liposomes (Friedmann (1989), supra; Brigham, et al. (1989) Am. J. Med. Sci., 298:278-281; Nabel, et al. (1990) Science, 249:1285-1288; Hazinski, et al. (1991) Am. J. Resp. Cell Molec. Biol., 4:206-209; and Wang and Huang (1987) Proc. Natl. Acad. Sci. (USA), 84:7851-7855); coupled to ligand-specific, cation-based transport systems (Wu and Wu (1988) J. Biol. Chem., 263:14621-14624) or the use of naked DNA, expression vectors (Nabel et al. (1990), supra); Wolff et al. (1990) Science, 247:1465-1468). Direct injection of transgenes into tissue produces only localized expression (Rosenfeld (1992) supra); Rosenfeld et al. (1991) supra; Brigham et al. (1989) supra; Nabel (1990) supra; and Hazinski et al. (1991) supra). The Brigham et al. group (Am. J. Med. Sci. (1989) 298:278-281 and Clinical Research (1991) 39 (abstract)) have reported in vivo transfection only of lungs of mice following either intravenous or intratracheal administration of a DNA liposome complex. An example of a review article of human gene therapy procedures is: Anderson, Science (1992) 256:808-813.

The antisense molecules of the invention encompass any pharmaceutically acceptable salts, esters, or salts of such esters, or any other compound which, upon administration to an animal including a human, is capable of providing (directly or indirectly) the biologically active metabolite or residue thereof. Accordingly, for example, the disclosure is also drawn to prodrugs and pharmaceutically acceptable salts of the compounds of the invention, pharmaceutically acceptable salts of such pro-drugs, and other bioequivalents.

The term "pharmaceutically acceptable salts" refers to physiologically and pharmaceutically acceptable salts of the compounds of the invention: i.e., salts that retain the desired biological activity of the parent compound and do not impart undesired toxicological effects thereto.

For oligonucleotides, preferred examples of pharmaceutically acceptable salts include but are not limited to (a) salts formed with cations such as sodium, potassium, ammonium, magnesium, calcium, polyamines such as spermine and spermidine, etc.; (b) acid addition salts formed with inorganic acids, for example hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid and the like; (c) salts formed with organic acids such as, for example, acetic acid, oxalic acid, tartaric acid, succinic acid, maleic acid, fumaric acid, gluconic acid, citric acid, malic acid, ascorbic acid, benzoic acid, tannic acid, palmitic acid, alginic acid, polyglutamic acid, naphthalenesulfonic acid, methanesulfonic acid, p-toluenesulfonic acid, naphthalenedisulfonic acid, polygalacturonic acid, and the like; and (d) salts formed from elemental anions such as chlorine, bromine, and iodine. The pharmaceutical compositions of the present invention may be administered in a number of ways depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical (including ophthalmic and to mucous membranes including rectal delivery), pulmonary, e.g., by inhalation or insufflation of powders or aerosols, (including by nebulizer, intratracheal, intranasal, epidermal and transdermal), oral or parenteral. Parenteral administration includes intravenous, intra-arterial, subcutaneous, intra-

US 8,455,636 B2

31

peritoneal or intramuscular injection or infusion; or intracranial, e.g., intrathecal or intraventricular, administration. Oligonucleotides with at least one 2'-O-methoxyethyl modification are believed to be particularly useful for oral administration.

The pharmaceutical formulations of the present invention, which may conveniently be presented in unit dosage form, may be prepared according to conventional techniques well known in the pharmaceutical industry. Such techniques include the step of bringing into association the active ingredients with the pharmaceutical carrier(s) or excipient(s). In general the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

Kits of the Invention

The invention also provides kits for treatment of a patient with a genetic disease which kit comprises at least an antisense molecule, packaged in a suitable container, together with instructions for its use.

In a preferred embodiment, the kits will contain at least one antisense molecule as shown in Table 1A, or a cocktail of antisense molecules as shown in Table 1B or a "weasel" compound as shown in Table 1C. The kits may also contain peripheral reagents such as buffers, stabilizers, etc.

Those of ordinary skill in the field should appreciate that applications of the above method has wide application for identifying antisense molecules suitable for use in the treatment of many other diseases.

EXAMPLES

The following Examples serve to more fully describe the manner of using the above-described invention, as well as to set forth the best modes contemplated for carrying out various aspects of the invention. It is understood that these Examples in no way serve to limit the true scope of this invention, but rather are presented for illustrative purposes. The references cited herein are expressly incorporated by reference.

Methods of molecular cloning, immunology and protein chemistry, which are not explicitly described in the following examples, are reported in the literature and are known by those skilled in the art. General texts that described conventional molecular biology, microbiology, and recombinant DNA techniques within the skill of the art, included, for example: Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989); Glover ed., *DNA Cloning: A Practical Approach*, Volumes I and II, MRL Press, Ltd., Oxford, U.K. (1985); and Ausubel, F., Brent, R., Kingston, R. E., Moore, D. D., Seidman, J. G., Smith, J. A., Struhl, K. *Current Protocols in Molecular Biology*. Greene Publishing Associates/Wiley Intersciences, New York (2002).

Determining Induced Exon Skipping in Human Muscle Cells Attempts by the inventors to develop a rational approach in antisense molecules design were not completely successful as there did not appear to be a consistent trend that could be applied to all exons. As such, the identification of the most effective and therefore most therapeutic antisense molecules compounds has been the result of empirical studies.

These empirical studies involved the use of computer programs to identify motifs potentially involved in the splicing process. Other computer programs were also used to identify regions of the pre-mRNA which may not have had extensive secondary structure and therefore potential sites for annealing of antisense molecules. Neither of these approaches proved

32

completely reliable in designing antisense oligonucleotides for reliable and efficient induction of exon skipping.

Annealing sites on the human dystrophin pre-mRNA were selected for examination, initially based upon known or predicted motifs or regions involved in splicing. 2OMe antisense oligonucleotides were designed to be complementary to the target sequences under investigation and were synthesised on an Expedite 8909 Nucleic Acid Synthesiser. Upon completion of synthesis, the oligonucleotides were cleaved from the support column and de-protected in ammonium hydroxide before being desalted. The quality of the oligonucleotide synthesis was monitored by the intensity of the trityl signals upon each deprotection step during the synthesis as detected in the synthesis log. The concentration of the antisense oligonucleotide was estimated by measuring the absorbance of a diluted aliquot at 260 nm.

Specified amounts of the antisense molecules were then tested for their ability to induce exon skipping in an in vitro assay, as described below.

Briefly, normal primary myoblast cultures were prepared from human muscle biopsies obtained after informed consent. The cells were propagated and allowed to differentiate into myotubes using standard culturing techniques. The cells were then transfected with the antisense oligonucleotides by delivery of the oligonucleotides to the cells as cationic lipoplexes, mixtures of antisense molecules or cationic liposome preparations.

The cells were then allowed to grow for another 24 hours, after which total RNA was extracted and molecular analysis commenced. Reverse transcriptase amplification (RT-PCR) was undertaken to study the targeted regions of the dystrophin pre-mRNA or induced exonic re-arrangements.

For example, in the testing of an antisense molecule for inducing exon 19 skipping the RT-PCR test scanned several exons to detect involvement of any adjacent exons. For example, when inducing skipping of exon 19, RT-PCR was carried out with primers that amplified across exons 17 and 21. Amplifications of even larger products in this area (i.e. exons 13-26) were also carried out to ensure that there was minimal amplification bias for the shorter induced skipped transcript. Shorter or exon skipped products tend to be amplified more efficiently and may bias the estimated of the normal and induced transcript.

The sizes of the amplification reaction products were estimated on an agarose gel and compared against appropriate size standards. The final confirmation of identity of these products was carried out by direct DNA sequencing to establish that the correct or expected exon junctions have been maintained.

Once efficient exon skipping had been induced with one antisense molecule, subsequent overlapping antisense molecules may be synthesized and then evaluated in the assay as described above. Our definition of an efficient antisense molecule is one that induces strong and sustained exon skipping at transfection concentrations in the order of 300 nM or less. Antisense Oligonucleotides Directed at Exon 8

Antisense oligonucleotides directed at exon 8 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

FIG. 3 shows differing efficiencies of two antisense molecules directed at exon 8 acceptor splice site. H8A(-06+18) [SEQ ID NO:1], which anneals to the last 6 bases of intron 7 and the first 18 bases of exon 8, induces substantial exon 8 and 9 skipping when delivered into cells at a concentration of 20 nM. The shorter antisense molecule, H8A(-06+14) [SEQ ID NO: 4] was only able to induce exon 8 and 9 skipping at 300

US 8,455,636 B2

33

nM, a concentration some 15 fold higher than H8A(-06+18), which is the preferred antisense molecule.

This data shows that some particular antisense molecules induce efficient exon skipping while another antisense molecule, which targets a near-by or overlapping region, can be much less efficient. Titration studies show one compound is able to induce targeted exon skipping at 20 nM while the less efficient antisense molecules only induced exon skipping at concentrations of 300 nM and above. Therefore, we have shown that targeting of the antisense molecules to motifs involved in the splicing process plays a crucial role in the overall efficacy of that compound.

Efficacy refers to the ability to induce consistent skipping of a target exon. However, sometimes skipping of the target exons is consistently associated with a flanking exon. That is, we have found that the splicing of some exons is tightly linked. For example, in targeting exon 23 in the mouse model of muscular dystrophy with antisense molecules directed at the donor site of that exon, dystrophin transcripts missing exons 22 and 23 are frequently detected. As another example, when using an antisense molecule directed to exon 8 of the human dystrophin gene, all induced transcripts are missing both exons 8 and 9. Dystrophin transcripts missing only exon 8 are not observed.

Table 2 below discloses antisense molecule sequences that induce exon 8 (and 9) skipping.

TABLE 2

(SEQ ID NOS 1-5, respectively, in order of appearance)		
Antisense Oligonucleotide name	Sequence	Ability to induce skipping
H8A(-06 + 18)	5'-GAU AGG UGG UAU CAA CAU CUG UAA	Very strong to 20 nM
H8A (-03 + 18)	5'-GAU AGG UGG UAU CAA CAU CUG	Very strong skipping to 40 nM
H8A(-07 + 18)	5'-GAU AGG UGG UAU CAA CAU CUG UAA G	Strong skipping to 40 nM
H8A(-06 + 14)	5'-GGU GGU AUC AAC AUC UGU AA	Skipping to 300 nM
H8A(-10 + 10)	5'-GUA UCA ACA UCU GUA AGC AC	Patchy/weak skipping to 100 nm

Table 2 (SEQ ID NOS 1-5, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 7

Antisense oligonucleotides directed at exon 7 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

FIG. 4 shows the preferred antisense molecule, H7A(+45+67) [SEQ ID NO: 6], and another antisense molecule, H7A(+2+26) [SEQ ID NO: 7], inducing exon 7 skipping. Nested amplification products span exons 3 to 9. Additional products above the induced transcript missing exon 7 arise from amplification from carry-over outer primers from the RT-PCR as well as heteroduplex formation.

Table 3 below discloses antisense molecule sequences for induced exon 7 skipping.

34

TABLE 3

(SEQ ID NOS 6-9, respectively, in order of appearance)		
Antisense Oligonucleotide name	Sequence	Ability to induce skipping
H7A(+45 + 67)	5'-UGC AUG UUC CAG UCG UUG UGU GG	Strong skipping to 20 nM
H7A(+02 + 26)	5'-CAC UAU UCC AGU CAA AUA GGU CUG G	Weak skipping at 100 nM
H7D(+15 - 10)	5'-AUU UAC CAA CCU UCA GGA UCG AGU A	Weak skipping to 300 nM
H7A(-18 + 03)	5'-GGC CUA AAA CAC AUA	Weak skipping to 300 nM

Table 3 (SEQ ID NOS 6-9, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 6

Antisense oligonucleotides directed at exon 6 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

FIG. 5 shows an example of two non-preferred antisense molecules inducing very low levels of exon 6 skipping in cultured human cells. Targeting this exon for specific removal was first undertaken during a study of the canine model using the oligonucleotides as listed in Table 4, below. Some of the human specific oligonucleotides were also evaluated, as shown in FIG. 5. In this example, both antisense molecules target the donor splice site and only induced low levels of exon 6 skipping. Both H6D(+4-21) [SEQ ID NO: 17] and H6D(+18-4) [SEQ ID NO: 18] would be regarded as non-preferred antisense molecules.

One antisense oligonucleotide that induced very efficient exon 6 skipping in the canine model, C6A(+69+91) [SEQ ID NO: 14], would anneal perfectly to the corresponding region in human dystrophin exon 6. This compound was evaluated, found to be highly efficient at inducing skipping of that target exon, as shown in FIG. 6 and is regarded as the preferred compound for induced exon 6 skipping. Table 4 below discloses antisense molecule sequences for induced exon 6 skipping.

TABLE 4

(SEQ ID NOS 10-18, respectively, in order of appearance)		
Antisense Oligo name	Sequence	Ability to induce skipping
C6A(-10 + 10)	5' CAU UUU UGA ACA UGU GG	No skipping
C6A(-14 + 06)	5' UUU GAC CUA GUG GAA AG	No skipping
C6A(-14 + 12)	5' UAC AUU UUU GAC CUA CAU GUG GAA AG	No skipping
C6A(-13 + 09)	5' AUU UUU GAC CUA CAU GGG AAA G	No skipping
CH6A(+69 + 91)	5' UAC GAG UUG AUU GUC GGA CCC AG	Strong skipping to 20 nM
C6D(+12 - 13)	5' GUG GUC UCC UUA CCU AUG ACU GUG G	Weak skipping at 300 nM

US 8,455,636 B2

35

TABLE 4-continued

(SEQ ID NOS 10-18, respectively, in order of appearance)		
Antisense Oligo name	Sequence	Ability to induce skipping
C6D(+06 - 11)	5' GGU CUC CUU ACC UAU GA	No skipping
H6D(+04 - 21)	5' UGU CUC AGU AAU CUU CUU ACC UAU	Weak skipping to 50 nM
H6D(+18 - 04)	5' UCU UAC CUA CUA UGG AUG AGA	Very weak skipping to 300 nM

Table 4 (SEQ ID NOS 10-18, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 4

Antisense oligonucleotides directed at exon 4 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

FIG. 7 shows an example of a preferred antisense molecule inducing skipping of exon 4 skipping in cultured human cells. In this example, one preferred antisense compound, H4A(+13+32) [SEQ ID NO:19], which targeted a presumed exonic splicing enhancer induced efficient exon skipping at a concentration of 20 nM while other non-preferred antisense oligonucleotides failed to induce even low levels of exon 4 skipping. Another preferred antisense molecule inducing skipping of exon 4 was H4A(+11+40) [SEQ ID NO:22], which induced efficient exon skipping at a concentration of 20 nM.

Table 5 below discloses antisense molecule sequences for inducing exon 4 skipping.

TABLE 5

(SEQ ID NOS 19, 22, 20, and 21, respectively, in order of appearance)		
Antisense Oligonucleotide name	Sequence	Ability to induce skipping
H4A(+13 + 32)	5' GCA UGA ACU CUU GUG GAU CC	Skipping to 20 nM
H4A(+11 + 40)	5'UGU UCA GGG CAU GAA CUC UUG UGG AUC CUU	Skipping to 20 nM
H4D(+04 - 16)	5' CCA GGG UAC UAC UUA CAU UA	No skipping
H4D(-24 - 44)	5' AUC GUG UGU CAC AGC AUC CAG	No skipping

Table 5 (SEQ ID NOS 19, 22, 20, and 21, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 3

Antisense oligonucleotides directed at exon 3 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

H3A(+30+60) [SEQ ID NO:23] induced substantial exon 3 skipping when delivered into cells at a concentration of 20 nM to 600 nM. The antisense molecule, H3A(+35+65) [SEQ ID NO: 24] induced exon skipping at 300 nM.

36

Table 6 below discloses antisense molecule sequences that induce exon 3 skipping.

TABLE 6

(SEQ ID NOS 23-30, respectively, in order of appearance)		
Antisense Oligonucleotide name	Sequence	Ability to induce skipping
H3A(+30 + 60)	UAG GAG GCG CCU CCC AUC CUG UAG GUC ACU G	Moderate skipping to 20 to 600 nM
H3A(+35 + 65)	AGG UCU AGG AGG GCG CUC CCA UCC UGU AGG U	Working to 300 nM
H3A(+30 + 54)	GCG CCU CCC AUC CUG UAG GUC ACU G	Moderate 100-600 nM
H3D(+46 - 21)	CUU CGA GGA GGU CUA GGA GGC GCC UC	No skipping
H3A(+30 + 50)	CUC CCA UCC UGU AGG UCA CUG	Moderate 20-600 nM
H3D(+19 - 03)	UAC CAG UUU UUG CCC UGU CAG G	No skipping
H3A(-06 + 20)	UCA AUA UGC UGC UUUCCA AAC UGA AA	No skipping
H3A(+37 + 61)	CUA GGA GGC GCC UCC CAU CCU GUA G	No skipping

Table 6 (SEQ ID NOS 23-30, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 5

Antisense oligonucleotides directed at exon 5 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

H5A(+20+50) [SEQ ID NO:31] induces substantial exon 5 skipping when delivered into cells at a concentration of 100 nM. Table 7 below shows other antisense molecules tested. The majority of these antisense molecules were not as effective at exon skipping as H5A(+20+50). However, H5A(+15+45) [SEQ ID NO: 40] was able to induce exon 5 skipping at 300 nM.

Table 7 below discloses antisense molecule sequences that induce exon 5 skipping.

TABLE 7

(SEQ ID NOS 31-40, respectively, in order of appearance)		
Antisense Oligonucleotide name	Sequence	Ability to induce skipping
H5A(+20 + 50)	UUA UGA UUU CCA UCU ACG AUG UCA GUA CUU C	Working to 100 nM
H5D(+25 - 05)	CUU ACC UGC CAG UGG AGG AUU AUA UUC CAA A	No skipping
H5D(+10 - 15)	CAU CAG GAU UCU UAC CUG CCA GUG G	Inconsistent at 300 nM
H5A(+10 + 34)	CGA UGU CAG UAC UUC CAA UAU UCA C	Very weak

US 8,455,636 B2

37

TABLE 7-continued

(SEQ ID NOS 31-40, respectively, in order of appearance)		
Antisense Oligonucleotide name	Sequence	Ability to induce skipping
H5D(-04 - 21)	ACC AUU CAU CAG GAU UCU	No skipping
H5D(+16 - 02)	ACC UGC CAG UGG AGG AUU	No skipping
H5A(-07 + 20)	CCA AUA UUC ACU AAA UCA ACC UGU UAA	No skipping
H5D(+18 - 12)	CAG GAU UCU UAC CUG CCA GUG GAG GAU UAU	No skipping
H5A(+05 + 35)	ACG AUG UCA GUA CUU CCA AUA UUC ACU AAA U	No skipping
H5A(+15 + 45)	AUU UCC AUC UAC GAU GUC AGU ACU UCC AAU A	Working to 300 nM

Table 7 (SEQ ID NOS 31-40, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 10

Antisense oligonucleotides directed at exon 10 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

H10A(-05+16) [SEQ ID NO:41] induced substantial exon 10 skipping when delivered into cells. Table 8 below shows other antisense molecules tested. The antisense molecules ability to induce exon skipping was variable. Table 8 below discloses antisense molecule sequences that induce exon 10 skipping.

TABLE 8

(SEQ ID NOS 41-45, respectively, in order of appearance)		
Antisense Oligonucleotide name	Sequence	Ability to induce skipping
H10A(-05 + 16)	CAG GAG CUU CCA AAU GCU GCA	Not tested
H10A(-05 + 24)	CUU GUC UUC AGG AGC UUC CAA AUG CUG CA	Not tested
H10A(+98 + 119)	UCC UCA GCA GAA AGA AGC CAC G	Not tested
H10A(+130 + 149)	UUA GAA AUC UCU CCU UGU GC	No skipping
H10A(-33 - 14)	UAA AUU GGG UGU UAC ACA AU	No skipping

Table 8 (SEQ ID NOS 41-45, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 11

Antisense oligonucleotides directed at exon 11 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

FIG. 8B shows an example of H11A(+75+97) [SEQ ID NO:49] antisense molecule inducing exon 11 skipping in cultured human cells. H11A(+75+97) induced substantial

38

exon 11 skipping when delivered into cells at a concentration of 5 nM. Table 9 below shows other antisense molecules tested. The antisense molecules ability to induce exon skipping was observed at 100 nM.

TABLE 9

(SEQ ID NOS 46-49 and 49, respectively, in order of appearance)		
Antisense Oligonucleotide name	Sequence	Ability to induce skipping
H11D(+26 + 49)	CCC UGA GGC AUU CCC AUC UUG AAU	Skipping at 100 nM
H11D(+11 - 09)	AGG ACU UAC UUG CUU UGU UU	Skipping at 100 nM
H11A(+118 + 140)	CUU GAA UUU AGG AGA UUC AUC UG	Skipping at 100 nM
H11A(+75 + 97)	CAU CUU CUG AUA AUU UUC CUG UU	Skipping at 100 nM
H11A(+75 + 97)	CAU CUU CUG AUA AUU UUC CUG UU	Skipping at 5 nM

Antisense Oligonucleotides Directed at Exon 12

Antisense oligonucleotides directed at exon 12 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

H12A(+52+75) [SEQ ID NO:50] induced substantial exon 12 skipping when delivered into cells at a concentration of 5 nM, as shown in FIG. 8A. Table 10 below shows other antisense molecules tested at a concentration range of 5, 25, 50, 100, 200 and 300 nM. The antisense molecules ability to induce exon skipping was variable.

TABLE 10

(SEQ ID NOS 50-52, respectively, in order of appearance)		
Antisense Oligonucleotide name	Sequence	Ability to induce skipping
H12A(+52 + 75)	UCU UCU GUU UUU GUU AGC CAG UCA	Skipping at 5 nM
H12A(-10 + 10)	UCU AUG UAA ACU GAA AAU UU	Skipping at 100 nM
H12A(+11 + 30)	UUC UGG AGA UCC AUU AAA AC	No skipping

Table 10 (SEQ ID NOS 50-52, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 13

Antisense oligonucleotides directed at exon 13 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

H13A(+77+100) [SEQ ID NO:53] induced substantial exon 13 skipping when delivered into cells at a concentration of 5 nM. Table 11 below includes two other antisense molecules tested at a concentration range of 5, 25, 50, 100, 200 and 300 nM. These other antisense molecules were unable to induce exon skipping.

US 8,455,636 B2

39

TABLE 11

(SEQ ID NOS 53-55, respectively, in order of appearance)		
Antisense Oligonucleotide name	Sequence	Ability to induce skipping
H13A(+77 + 100)	CAG CAG UUG CGU GAU CUC CAC UAG	Skipping at 5 nM
H13A(+55 + 75)	UUC AUC AAC UAC CAC CAC CAU	No skipping
H13D(+06 - 19)	CUA AGC AAA AUA AUC UGA CCU UAA G	No skipping

Table 11 (SEQ ID NOS 53-55, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 14

Antisense oligonucleotides directed at exon 14 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

H14A(+37+64) [SEQ ID NO:56] induced weak exon 14 skipping when delivered into cells at a concentration of 100 nM. Table 12 below includes other antisense molecules tested at a concentration range of 5, 25, 50, 100, 200 and 300 nM. The other antisense molecules were unable to induce exon skipping at any of the concentrations tested.

TABLE 12

(SEQ ID NOS 56-62, respectively, in order of appearance)		
Antisense Oligonucleotide name	Sequence	Ability to induce skipping
H14A(+37 + 64)	CUU GUA AAA GAA CCC AGC GGU CUU CUG U	Skipping at 100 nM
H14A(+14 + 35)	CAU CUA CAG AUG UUU GCC CAU C	No skipping
H14A(+51 + 73)	GAA GGA UGU CUU GUA AAA GAA CC	No skipping
H14D(-02 + 18)	ACC UGU UCU UCA GUA AGA CG	No skipping
H14D(+14 - 10)	CAU GAC ACA CCU GUU CUU CAG UAA	No skipping
H14A(+61 + 80)	CAU UUG AGA AGG AUG UCU UG	No skipping
H14A(-12 + 12)	AUC UCC CAA UAC CUG GAG AAG AGA	No skipping

40

Table 12 (SEQ ID NOS 56-62, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 15

5 Antisense oligonucleotides directed at exon 15 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

10 H15A(-12+19) [SEQ ID NO:63] and H15A(+48+71) [SEQ ID NO:64] induced substantial exon 15 skipping when delivered into cells at a concentration of 10 nM, as shown in FIG. 9A. Table 13 below includes other antisense molecules tested at a concentration range of 5, 25, 50, 100, 200 and 300 nM. These other antisense molecules were unable to induce exon skipping at any of the concentrations tested.

TABLE 13

(SEQ ID NOS 63-65, 63, and 66, respectively, in order of appearance)		
Antisense Oligonucleotide name	Sequence	Ability to induce skipping
25 H15A(-12 + 19)	GCC AUG CAC UAA AAA GGC ACU GCA AGA CAU U	Skipping at 5 nM
H15A(+48 + 71)	UCU UUA AAG CCA GUU GUG UGA AUC	Skipping at 5 nM
30 H15A(+08 + 28)	UUU CUG AAA GCC AUG CAC UAA	No skipping
H15A(-12 + 19)	GCC AUG CAC UAA AAA GGC ACU GCA AGA CAU U	Skipping at 10 nM
35 H15D(+17 - 08)	GUA CAU ACG GCC AGU UUU UGA AGA C	No skipping

Antisense Oligonucleotides Directed at Exon 16

40 Antisense oligonucleotides directed at exon 16 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

45 H16A(-12+19) [SEQ ID NO:67] and H16A(-06+25) [SEQ ID NO:68] induced substantial exon 16 skipping when delivered into cells at a concentration of 10 nM, as shown in FIG. 9B. Table 14 below includes other antisense molecules tested. H16A(-06+19) [SEQ ID NO:69] and H16A(+87+109) [SEQ ID NO:70] were tested at a concentration range of 5, 25, 50, 100, 200 and 300 nM. These two antisense molecules were able to induce exon skipping at 25 nM and 100 nM, respectively. Additional antisense molecules were tested at 100, 200 and 300 nM and did not result in any exon skipping.

TABLE 14

(SEQ ID NOS 67-78, respectively, in order of appearance)		
Antisense Oligonucleotide name	Sequence	Ability to induce skipping
H16A(-12 + 19)	CUA GAU CCG CUU UUA AAA CCU GUU AAA ACA A	Skipping at 5 nM

US 8,455,636 B2

41

TABLE 14-continued

(SEQ ID NOS 67-78, respectively, in order of appearance)		
Antisense Oligonucleotide name	Sequence	Ability to induce skipping
H16A(-06 + 25)	UCU UUU CUA GAU CCG CUU UUA AAA CCU GUU A	Skipping at 5 nM
H16A(-06 + 19)	CUA GAU CCG CUU UUA AAA CCU GUU A	Skipping at 25 nM
H16A(+87 + 109)	CCG UCU UCU GGG UCA CUG ACU UA	Skipping at 100 nM
H16A(-07 + 19)	CUA GAU CCG CUU UUA AAA CCU GUU AA	No skipping
H16A(-07 + 13)	CCG CUU UUA AAA CCU GUU AA	No skipping
H16A(+12 + 37)	UGG AUU GCU UUU UCU UUU CUA GAU CC	No skipping
H16A(+92 + 116)	CAU GCU UCC GUC UUC UGG GUC ACU G	No skipping
H16A(+45 + 67)	G AUC UUG UUU GAG UGA AUA CAG U	No skipping
H16A(+105 + 126)	GUU AUC CAG CCA UGC UUC CGU C	No skipping
H16D(+05 - 20)	UGA UAA UUG GUA UCA CUA ACC UGU G	No skipping
H16D(+12 - 11)	GUA UCA CUA ACC UGU GCU GUA C	No skipping

42

Table 14 (SEQ ID NOS 67-78, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 19

Antisense oligonucleotides directed at exon 19 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

H19A(+35+65) [SEQ ID NO:79] induced substantial exon 19 skipping when delivered into cells at a concentration of 10 nM. This antisense molecule also showed very strong exon skipping at concentrations of 25, 50, 100, 300 and 600 nM.

FIG. 10 illustrates exon 19 and 20 skipping using a “cocktail” of antisense oligonucleotides, as tested using gel electrophoresis. It is interesting to note that it was not easy to induce exon 20 skipping using single antisense oligonucleotides H20A(+44+71) [SEQ ID NO:81] or H20A(+149+170) [SEQ ID NO:82], as illustrated in sections 2 and 3 of the gel shown in FIG. 10. Whereas, a “cocktail” of antisense oligonucleotides was more efficient as can be seen in section 4 of FIG. 10 using a “cocktail” of antisense oligonucleotides H20A(+44+71) and H20A(+149+170). When the cocktail was used to target exon 19, skipping was even stronger (see section 5, FIG. 10).

FIG. 11 illustrates gel electrophoresis results of exon 19/20 skipping using “weasels” The “weasels” were effective in skipping exons 19 and 20 at concentrations of 25, 50, 100, 300 and 600 nM. A further “weasel” sequence is shown in the last row of Table 3C. This compound should give good results.

Antisense Oligonucleotides Directed at Exon 20

Antisense oligonucleotides directed at exon 20 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

None of the antisense oligonucleotides tested induced exon 20 skipping when delivered into cells at a concentration of 10, 25, 50, 300 or 600 nM (see Table 15). Antisense molecules H20A(-11+17) [SEQ ID NO:86] and H20D(+08-20) [SEQ ID NO:87] are yet to be tested.

However, a combination or “cocktail” of H20A(+44+71) [SEQ ID NO: 81] and H20A(+147+168) [SEQ ID NO:82] in a ratio of 1:1, exhibited very strong exon skipping at a concentration of 100 nM and 600 nM. Further, a combination of antisense molecules H19A(+35+65) [SEQ ID NO:80], H20A(+44+71) [SEQ ID NO:81] and H20A(+147+168) [SEQ ID NO:82] in a ratio of 2:1:1, induced very strong exon skipping at a concentration ranging from 10 nM to 600 nM.

TABLE 15

(SEQ ID NOS 81-87, 81-82, and 80-82, respectively, in order of appearance)		
Antisense Oligonucleotide name	Sequence	Ability to induce skipping
H20A(+44 + 71)	CUG GCA GAA UUC GAU CCA CCG GCU GUU C	No skipping
H20A(+147 + 168)	CAG CAG UAG UUG UCA UCU GCU C	No skipping
H20A(+185 + 203)	UGA UGG GGU GGU GGG UUG G	No skipping
H20A(-08 + 17)	AUC UGC AUU AAC ACC CUC UAG AAA G	No skipping

US 8,455,636 B2

43

TABLE 15-continued

(SEQ ID NOS 81-87, 81-82, and 80-82, respectively, in order of appearance)		
Antisense Oligonucleotide name	Sequence	Ability to induce skipping
H20A(+30 + 53)	CCG GCU GUU CAG UUG UUC UGA GGC	No skipping
H20A(-11 + 17)	AUC UGC AUU AAC ACC CUC UAG AAA GAA A	Not tested yet
H20D(+08 - 20)	GAA GGA GAA GAG AUU CUU ACC UUA CAA A	Not tested yet
H20A(+44 + 71) & H20A(+147 + 168)	CUG GCA GAA UUC GAU CCA CCG GCU GUU C CAG CAG UAG UUG UCA UCU GCU C	Very strong skipping
H19A(+35 + 65); H20A(+44 + 71); H20A(+147 + 168)	GCC UGA GCU GAU CUG CUG GCA UCU UGC AGU U CUG GCA GAA UUC GAU CCA CCG GCU GUU C CAG CAG UAG UUG UCA UCU GCU C	Very strong skipping

44

Antisense Oligonucleotides Directed at Exon 21

Antisense oligonucleotides directed at exon 21 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

H21A(+85+108) [SEQ ID NO:92] and H21A(+85+106) [SEQ ID NO:91] induced exon 21 skipping when delivered into cells at a concentration of 50 nM. Table 16 below includes other antisense molecules tested at a concentration range of 5, 25, 50, 100, 200 and 300 nM. These antisense molecules showed a variable ability to induce exon skipping

TABLE 16

(SEQ ID NOS 90-94, respectively, in order of appearance)		
Antisense Oligonucleotide name	Sequence	Ability to induce skipping
H21A(-06 + 16)	GCC GGU UGA CUU CAU CCU GUG C	Skips at 600 nM
H21A(+85 + 106)	CUG CAU CCA GGA ACA UGG GUC C	Skips at 50 nM
H21A(+85 + 108)	GUC UGC AUC CAG GAA CAU GGG UC	Skips at 50 nM
H21A(+08 + 31)	GUU GAA GAU CUG AUA GCC GGU UGA	Skips faintly to

TABLE 16-continued

(SEQ ID NOS 90-94, respectively, in order of appearance)		
Antisense Oligonucleotide name	Sequence	Ability to induce skipping
H21D(+18 - 07)	UAC UUA CUG UCU GUA GCU CUU UCU	No skipping

Table 16 (SEQ ID NOS 90-94, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 22

Antisense oligonucleotides directed at exon 22 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

FIG. 12 illustrates differing efficiencies of two antisense molecules directed at exon 22 acceptor splice site. H22A(+125+106) [SEQ ID NO:96] and H22A(+80+101) [SEQ ID NO: 98] induce strong exon 22 skipping from 50 nM to 600 nM concentration.

H22A(+125+146) [SEQ ID NO:96] and H22A(+80+101) [SEQ ID NO:98] induced exon 22 skipping when delivered into cells at a concentration of 50 nM. Table 17 below shows other antisense molecules tested at a concentration range of 50, 100, 300 and 600 nM. These antisense molecules showed a variable ability to induce exon skipping.

TABLE 17

(SEQ ID NOS 95-99, respectively, in order of appearance)		
Antisense oligonucleotide name	Sequence	Ability to induce skipping
H22A(+22 + 45)	CAC UCA UGG UCU CCU GAU AGC GCA	No skipping
H22A(+125 + 146)	CUG CAA UUC CCC GAG UCU CUG C	Skipping to 50 nM
H22A(+47 + 69)	ACU GCU GGA CCC AUG UCC UGA UG	Skipping to 300 nM
H22A(+80 + 101)	CUA AGU UGA GGU AUG GAG AGU	Skipping to 50 nM
H22D(+13 - 11)	UAU UCA CAG ACC UGC AAU UCC CC	No skipping

US 8,455,636 B2

45

Table 17 (SEQ ID NOS 95-99, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 23

Antisense oligonucleotides directed at exon 23 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

Table 18 below shows antisense molecules tested at a concentration range of 25, 50, 100, 300 and 600 nM. These antisense molecules showed no ability to induce exon skipping or are yet to be tested.

TABLE 18

(SEQ ID NOS 100-102, respectively, in order of appearance)		
Antisense oligonucleotide name	Sequence	Ability to induce skipping
H23A(+34 + 59)	ACA GUG GUG CUG AGA UAG UAU AGG CC	No skipping
H23A(+18 + 39)	UAG GCC ACU UUG UUG CUC UUG C	No Skipping
H23A(+72 + 90)	UUC AGA GGG CGC UUU CUU C	No Skipping

Table 18 (SEQ ID NOS 100-102, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 24

Antisense oligonucleotides directed at exon 24 were prepared using similar methods as described above. Table 19 below outlines the antisense oligonucleotides directed at exon 24 that are yet to be tested for their ability to induce exon 24 skipping.

TABLE 19

(SEQ ID NOS 103-104, respectively, in order of appearance)		
Antisense oligonucleotide name	Sequence	Ability to induce skipping
H24A(+48 + 70)	GGG CAG GCC AUU CCU CCU UCA GA	Needs testing
H24A(-02 + 22)	UCU UCA GGG UUU GUA UGU GAU UCU	Needs testing

Table 19 (SEQ ID NOS 103-104, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 25

Antisense oligonucleotides directed at exon 25 were prepared using similar methods as described above. Table 20 below shows the antisense oligonucleotides directed at exon 25 that are yet to be tested for their ability to induce exon 25 skipping.

46

TABLE 20

(SEQ ID NOS 105-107, respectively, in order of appearance)		
Antisense oligonucleotide name	Sequence	Ability to induce skipping
H25A(+9 + 36)	CUG GGC UGA AUU GUC UGA AUA UCA CUG	Needs testing
H25A(+131 + 156)	CUG UUG GCA CAU GUG AUC CCA CUG AG	Needs testing
H25D(+16 - 08)	GUC UAU ACC UGU UGG CAC AUG UGA	Needs testing

Table 20 (SEQ ID NOS 105-107, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 26

Antisense oligonucleotides directed at exon 26 were prepared using similar methods as described above. Table 21 below outlines the antisense oligonucleotides directed at exon 26 that are yet to be tested for their ability to induce exon 26 skipping.

TABLE 21

(SEQ ID NOS 108-110, respectively, in order of appearance)		
Antisense oligonucleotide name	Sequence	Ability to induce skipping
H26A(+132 + 156)	UGC UUU CUG UAA UUC AUC UGG AGU U	Needs testing
H26A(-07 + 19)	CCU CCU UUC UGG CAU AGA CCU UCC AC	Needs testing
H26A(+68 + 92)	UGU GUC AUC CAU UGC UGC AUC UCU G	Faint skipping at 600 nM

Table 21 (SEQ ID NOS 108-110, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 27

Antisense oligonucleotides directed at exon 27 were prepared using similar methods as described above. Table 22 below outlines the antisense oligonucleotides directed at exon 27 that are yet to be tested for their ability to induce exon 27 skipping.

TABLE 22

(SEQ ID NOS 111-113, respectively, in order of appearance)		
Antisense oligonucleotide name	Sequence	Ability to induce skipping
H27A(+82 + 106)	UUA AGG CCU CUU GUG CUA CAG GUG G	Needs testing
H27A(-4 + 19)	GGG CCU CUU CUU UAG CUC UCU GA	Faint skipping at 600 and 300 nM
H27D(+19 - 03)	GAC UUC CAA AGU CUU GCA UUU C	v. strong skipping at 600 and 300 nM

Table 22 (SEQ ID NOS 111-113, respectively, in order of appearance)

US 8,455,636 B2

47

Antisense Oligonucleotides Directed at Exon 28

Antisense oligonucleotides directed at exon 28 were prepared using similar methods as described above. Table 23 below outlines the antisense oligonucleotides directed at exon 28 that are yet to be tested for their ability to induce exon 28 skipping.

TABLE 23

(SEQ ID NOS 114-116, respectively, in order of appearance)		
Antisense oligonucleotide name	Sequence	Ability to induce skipping
H28A(-05 + 19)	GCC AAC AUG CCC AAA CUU CCU AAG	v. strong skipping at 600 and 300 nM
H28A(+99 + 124)	CAG AGA UUU CCU CAG CUC CGC CAG GA	Needs testing
H28D(+16 - 05)	CUU ACA UCU AGC ACC UCA GAG	v. strong skipping at 600 and 300 nM

Table 23 (SEQ ID NOS 114-116, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 29

Antisense oligonucleotides directed at exon 29 were prepared using similar methods as described above. Table 24 below outlines the antisense oligonucleotides directed at exon 29 that are yet to be tested for their ability to induce exon 29 skipping.

TABLE 24

(SEQ ID NOS 117-119, respectively, in order of appearance)		
Antisense oligonucleotide name	Sequence	Ability to induce skipping
H29A(+57 + 81)	UCC GCC AUC UGU UAG GGU CUG UGC C	Needs testing
H29A(+18 + 42)	AUU UGG GUU AUC CUC UGA AUG UCG C	v. strong skipping at 600 and 300 nM
H29D(+17 - 05)	CAU ACC UCU UCA UGU AGU UCC C	v. strong skipping at 600 and 300 nM

Table 24 (SEQ ID NOS 117-119, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 30

Antisense oligonucleotides directed at exon 30 were prepared using similar methods as described above. Table 25 below outlines the antisense oligonucleotides directed at exon 30 that are yet to be tested for their ability to induce exon 30 skipping.

48

TABLE 25

(SEQ ID NOS 120-122, respectively, in order of appearance)		
Antisense oligonucleotide name	Sequence	Ability to induce skipping
H30A(+122 + 147)	CAU UUG AGC UGC GUC CAC CUU GUC UG	Needs testing
H30A(+25 + 50)	UCC UGG GCA GAC UGG AUG CUC UGU	Very strong skipping at 600 and 300 nM.
H30D(+19 - 04)	UUG CCU GGG CUU CCU GAG GCA UU	Very strong skipping at 600 and 300 nM.

Table 25 (SEQ ID NOS 120-122, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 31

Antisense oligonucleotides directed at exon 31 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

FIG. 13 illustrates differing efficiencies of two antisense molecules directed at exon 31 acceptor splice site and a “cocktail” of exon 31 antisense oligonucleotides at varying concentrations. H31 D(+03-22) [SEQ ID NO:124] substantially induced exon 31 skipping when delivered into cells at a concentration of 20 nM. Table 26 below also includes other antisense molecules tested at a concentration of 100 and 300 nM. These antisense molecules showed a variable ability to induce exon skipping.

TABLE 26

(SEQ ID NOS 123-126, respectively, in order of appearance)		
Antisense oligonucleotide name	Sequence	Ability to induce skipping
H31D(+06 - 18)	UUC UGA AAU AAC AUA UAC CUG UGC	Skipping to 300 nM
H31D(+03 - 22)	UAG UUU CUG AAA UAA CAU AUA CCU G	Skipping to 20 nM
H31A(+05 + 25)	GAC UUG UCA AAU CAG AUU GGA	No skipping
H31D(+04 - 20)	GUU UCU GAA AUA ACA UAU ACC UGU	Skipping to 300 nM

Table 26 (SEQ ID NOS 123-126, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 32

Antisense oligonucleotides directed at exon 32 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

H32D(+04-16) [SEQ ID NO:127] and H32A(+49+73) [SEQ ID NO:130] induced exon 32 skipping when delivered into cells at a concentration of 300 nM. Table 27 below also shows other antisense molecules tested at a concentration of 100 and 300 nM. These antisense molecules did not show an ability to induce exon skipping.

US 8,455,636 B2

49

TABLE 27

(SEQ ID NOS 127-130, respectively, in order of appearance)		
Antisense oligonucleotide name	Sequence	Ability to induce skipping
H32D(+04 - 16)	CAC CAG AAA UAC AUA CCA CA	Skipping to 300 nM
H32A(+151 + 170)	CAA UGA UUU AGC UGU GAC UG	No skipping
H32A(+10 + 32)	CGA AAC UUC AUG GAG ACA UCU UG	No skipping
H32A(+49 + 73)	CUU GUA GAC GCU GCU CAA AAU UGG C	Skipping to 300 nM

Table 27 (SEQ ID NOS 127-130, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 33

Antisense oligonucleotides directed at exon 33 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

FIG. 14 shows differing efficiencies of two antisense molecules directed at exon 33 acceptor splice site. H33A(+64+88) [SEQ ID NO:134] substantially induced exon 33 skipping when delivered into cells at a concentration of 10 nM. Table 28 below includes other antisense molecules tested at a concentration of 100, 200 and 300 nM. These antisense molecules showed a variable ability to induce exon skipping.

TABLE 28

(SEQ ID NOS 131-134, respectively, in order of appearance)		
Antisense oligonucleotide name	Sequence	Ability to induce skipping
H33D(+09 - 11)	CAU GCA CAC ACC UUU GCU CC	No skipping
H33A(+53 + 76)	UCU GUA CAA UCU GAC GUC CAG UCU	Skipping to 200 nM
H33A(+30 + 56)	GUC UUU AUC ACC AUU UCC ACU UCA GAC	Skipping to 200 nM
H33A(+64 + 88)	CCG UCU GCU UUU UCU GUA CAA UCU G	Skipping to 10 nM

Table 28 (SEQ ID NOS 131-134, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 34

Antisense oligonucleotides directed at exon 34 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

Table 29 below includes antisense molecules tested at a concentration of 100 and 300 nM. These antisense molecules showed a variable ability to induce exon skipping.

50

TABLE 29

(SEQ ID NOS 135-141, respectively, in order of appearance)		
Antisense oligonucleotide name	Sequence	Ability to induce skipping
H34A(+83 + 104)	UCC AUA UCU GUA GCU GCC AGC C	No skipping
H34A(+143 + 165)	CCA GGC AAC UUC AGA AUC CAA AU	No skipping
H34A(-20 + 10)	UUU CUG UUA CCU GAA AAG AAU UAU AAU GAA	Not tested
H34A(+46 + 70)	CAU UCA UUU CCU UUC GCA UCU UAC G	Skipping to 300 nM
H34A(+95 + 120)	UGA UCU CUU UGU CAA UUC CAU AUC UG	Skipping to 300 nM
H34D(+10 - 20)	UUC AGU GAU AUA GGU UUU ACC UUU CCC CAG	Not tested
H34A(+72 + 96)	CUG UAG CUG CCA GCC AUU CUG UCA AG	No skipping

Table 29 (SEQ ID NOS 135-141, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 35

Antisense oligonucleotides directed at exon 35 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

FIG. 15 shows differing efficiencies of antisense molecules directed at exon 35 acceptor splice site. H35A(+24+43) [SEQ ID NO:144] substantially induced exon 35 skipping when delivered into cells at a concentration of 20 nM. Table 30 below also includes other antisense molecules tested at a concentration of 100 and 300 nM. These antisense molecules showed no ability to induce exon skipping.

TABLE 30

(SEQ ID NOS 142-144, respectively, in order of appearance)		
Antisense oligonucleotide name	Sequence	Ability to induce skipping
H35A(+141 + 161)	UCU UCU GCU CGG GAG GUG ACA	Skipping to 20 nM
H35A(+116 + 135)	CCA GUU ACU AUU CAG AAG AC	No skipping
H35A(+24 + 43)	UCU UCA GGU GCA CCU UCU GU	No skipping

Table 30 (SEQ ID NOS 142-144, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 36

Antisense oligonucleotides directed at exon 36 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

US 8,455,636 B2

51

Antisense molecule H36A(+26+50) [SEQ ID NO:145] induced exon 36 skipping when delivered into cells at a concentration of 300 nM, as shown in FIG. 16.

Antisense Oligonucleotides Directed at Exon 37

Antisense oligonucleotides directed at exon 37 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

FIG. 17 shows differing efficiencies of two antisense molecules directed at exon 37 acceptor splice site. H37A(+82+105) [SEQ ID NO:148] and H37A(+134+157) [SEQ ID NO:149] substantially induced exon 37 skipping when delivered into cells at a concentration of 10 nM. Table 31 below shows the antisense molecules tested.

TABLE 31

(SEQ ID NOS 147-149, respectively, in order of appearance)		
Antisense oligonucleotide name	Sequence	Ability to induce skipping
H37A(+26 + 50)	CGU GUA GAG UCC ACC UUU GGG CGU A	No skipping
H37A(+82 + 105)	UAC UAA UUU CCU GCA GUG GUC ACC	Skipping to 10 nM
H37A(+134 + 157)	UUC UGU GUG AAA UGG CUG CAA AUC	Skipping to 10 nM

Table 31 (SEQ ID NOS 147-149, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 38

Antisense oligonucleotides directed at exon 38 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

FIG. 18 illustrates antisense molecule H38A(+88+112) [SEQ ID NO:152], directed at exon 38 acceptor splice site. H38A(+88+112) substantially induced exon 38 skipping when delivered into cells at a concentration of 10 nM. Table 32 below shows the antisense molecules tested and their ability to induce exon skipping.

TABLE 32

(SEQ ID NOS 150-152, respectively, in order of appearance)		
Antisense oligonucleotide name	Sequence	Ability to induce skipping
H38A(-01 + 19)	CCU UCA AAG GAA UGG AGG CC	No skipping
H38A(+59 + 83)	UGC UGA AUU UCA GCC UCC AGU GGU U	Skipping to 10 nM
H38A(+88 + 112)	UGA AGU CUU CCU CUU UCA GAU UCA C	Skipping to 10 nM

52

Table 32 (SEQ ID NOS 150-152, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 39

Antisense oligonucleotides directed at exon 39 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

H39A(+62+85) [SEQ ID NO:153] induced exon 39 skipping when delivered into cells at a concentration of 100 nM. Table 33 below shows the antisense molecules tested and their ability to induce exon skipping.

TABLE 33

(SEQ ID NOS 153-156, respectively, in order of appearance)		
Antisense oligonucleotide name	Sequence	Ability to induce skipping
H39A(+62 + 85)	CUG GCU UUC UCU CAU CUG UGA UUC	Skipping to 100 nM
H39A(+39 + 58)	GUU GUA AGU UGU CUC CUC UU	No skipping
H39A(+102 + 121)	UUG UCU GUA ACA GCU GCU GU	No skipping
H39D(+10 - 10)	GCU CUA AUA CCU UGA GAG CA	Skipping to 300 nM

Table 33 (SEQ ID NOS 153-156, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 40

Antisense oligonucleotides directed at exon 40 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

FIG. 19 illustrates antisense molecule H40A(-05+17) [SEQ ID NO:157] directed at exon 40 acceptor splice site. H40A(-05+17) and H40A(+129+153) [SEQ ID NO:158] both substantially induced exon 40 skipping when delivered into cells at a concentration of 5 nM.

Antisense Oligonucleotides Directed at Exon 42

Antisense oligonucleotides directed at exon 42 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

FIG. 20 illustrates antisense molecule H42A(-04+23) [SEQ ID NO:159], directed at exon 42 acceptor splice site. H42A(-4+23) and H42D(+19-02) [SEQ ID NO:161] both induced exon 42 skipping when delivered into cells at a concentration of 5 nM. Table 34 below shows the antisense molecules tested and their ability to induce exon 42 skipping.

TABLE 34

(SEQ ID NOS 159-160, respectively, in order of appearance)		
Antisense oligonucleotide name	Sequence	Ability to induce skipping
H42A(-4 + 23)	AUC GUU UCU UCA CGG ACA GUG UGC UGG	Skipping to 5 nM
H42A(+86 + 109)	GGG CUU GUG AGA CAU GAG UGA UUU	Skipping to 100 nM
H42D(+19 - 02)	A CCU UCA GAG GAC UCC UCU UGC	Skipping to 5 nM

US 8,455,636 B2

53

Table 34 (SEQ ID NOS 159-160, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 43

Antisense oligonucleotides directed at exon 43 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

H43A(+101+120) [SEQ ID NO:163] induced exon 43 skipping when delivered into cells at a concentration of 25 nM. Table 35 below includes the antisense molecules tested and their ability to induce exon 43 skipping.

TABLE 35

(SEQ ID NOS 162-164, respectively, in order of appearance)		
Antisense oligonucleotide name	Sequence	Ability to induce skipping
H43D(+10 - 15)	UAU GUG UUA CCU ACC CUU GUC GGU C	Skipping to 100 nM
H43A(+101 + 120)	GGA GAG AGC UUC CUG UAG CU	Skipping to 25 nM
H43A(+78 + 100)	UCA CCC UUU CCA CAG GCG UUG CA	Skipping to 200 nM

Table 35 (SEQ ID NOS 162-164, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 44

Antisense oligonucleotides directed at exon 44 were prepared using similar methods as described above. Testing for the ability of these antisense molecules to induce exon 44 skipping is still in progress. The antisense molecules under review are shown as SEQ ID Nos: 165 to 167 in Table 1A.

Antisense Oligonucleotides Directed at Exon 45

Antisense oligonucleotides directed at exon 45 were prepared using similar methods as described above. Testing for the ability of these antisense molecules to induce exon 45 skipping is still in progress. The antisense molecules under review are shown as SEQ ID Nos: 207 to 211 in Table 1A.

Antisense Oligonucleotides Directed at Exon 46

Antisense oligonucleotides directed at exon 46 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

FIG. 21 illustrates the efficiency of one antisense molecule directed at exon 46 acceptor splice site. Antisense oligonucleotide H46A(+86+115) [SEQ ID NO:203] showed very strong ability to induce exon 46 skipping. Table 36 below includes antisense molecules tested. These antisense molecules showed varying ability to induce exon 46 skipping.

TABLE 36

(SEQ ID NOS 168-169 and 203-206, respectively, in order of appearance)		
Antisense oligonucleotide name	Sequence	Ability to induce skipping
H46D(+16 - 04)	UUA CCU UGA CUU GCU CAA GC	No skipping
H46A(+90 + 109)	UCC AGG UUC AAG UGG GAU AC	No skipping

54

TABLE 36-continued

(SEQ ID NOS 168-169 and 203-206, respectively, in order of appearance)

Antisense oligonucleotide name	Sequence	Ability to induce skipping
H46A(+86 + 115)	CUC UUU UCC AGG UUC AAG UGG GAU ACU AGC	Good skipping to 100 nM
H46A(+107 + 137)	CAA GCU UUU CUU UUA GUU GCU GCU CUU UUC C	Good skipping to 100 nM
H46A(-10 + 20)	UAU UCU UUU GUU CUU CUA GCC UGG AGA AAG	Weak skipping
H46A(+50 + 77)	CUG CUU CCU CCA ACC AUA AAA CAA AUU C	Weak skipping

Table 36 (SEQ ID NOS 168-169 and 203-206, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 47

Antisense oligonucleotides directed at exon 47 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

H47A(+76+100) [SEQ ID NO:170] and H47A(-09+12) [SEQ ID NO:172] both induced exon 47 skipping when delivered into cells at a concentration of 200 nM. H47D(+25-02) [SEQ ID NO: 171] is yet to be prepared and tested.

Antisense Oligonucleotides Directed at Exon 50

Antisense oligonucleotides directed at exon 50 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

Antisense oligonucleotide molecule H50A(+02+30) [SEQ ID NO: 173] was a strong inducer of exon skipping. Further, H50A(+07+33) [SEQ ID NO:174] and H50D(+07-18) [SEQ ID NO:175] both induced exon 50 skipping when delivered into cells at a concentration of 100 nM.

Antisense Oligonucleotides Directed at Exon 51

Antisense oligonucleotides directed at exon 51 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

FIG. 22 illustrates differing efficiencies of two antisense molecules directed at exon 51 acceptor splice site. Antisense oligonucleotide H51A(+66+90) [SEQ ID NO:180] showed the stronger ability to induce exon 51 skipping. Table 37 below includes antisense molecules tested at a concentration range of 25, 50, 100, 300 and 600 nM. These antisense molecules showed varying ability to induce exon skipping. The strongest inducers of exon skipping were antisense oligonucleotide H51A(+61+90) [SEQ ID NO: 179] and H51A(+66+95) [SEQ ID NO: 179].

US 8,455,636 B2

55

TABLE 37

(SEQ ID NOS 176-185, respectively, in order of appearance)				
Antisense oligonucleotide name	Sequence	Ability to induce skipping		
H51A(-01 + 25)	ACC AGA GUA ACA GUC UGA GUA GGA GC	Faint skipping		
H51D(+16 - 07)	CUC AUA CCU UCU GCU UGA UGA UC	Skipping at 300 nM		
H51A(+111 + 134)	UUC UGU CCA AGC CCG GUU GAA AUC	Needs re-testing		
H51A(+61 + 90)	ACA UCA AGG AAG AUG GCA UUU CUA GUU UGG	Very strong skipping		
H51A(+66 + 90)	ACA UCA AGG AAG AUG GCA UUU CUA G	skipping		
H51A(+66 + 95)	CUC CAA CAU CAA GGA AGA UGG CAU UUC UAG	Very strong skipping		
H51D(+08 - 17)	AUC AUU UUU UCU CAU ACC UUC UGC U	No skipping		
H51A/D(+08 - 17) & (-15 + ?)	AUC AUU UUU UCU CAU ACC UUC UGC UAG GAG CUA AAA	No skipping		
H51A(+175 + 195)	CAC CCA CCA UCA CCC UCU GUG	No skipping		
H51A(+199 + 220)	AUC AUC UCG UUG AUA UCC UCA A	No skipping		

Antisense Oligonucleotides Directed at Exon 52

Antisense oligonucleotides directed at exon 52 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

FIG. 22 also shows differing efficiencies of four antisense molecules directed at exon 52 acceptor splice site. The most effective antisense oligonucleotide for inducing exon 52 skipping was H52A(+17+37) [SEQ ID NO:188].

Table 38 below shows antisense molecules tested at a concentration range of 50, 100, 300 and 600 nM. These antisense molecules showed varying ability to induce exon 50 skipping. Antisense molecules H52A(+12+41) [SEQ ID NO:187] and H52A(+17+37) [SEQ ID NO:188] showed the strongest exon 50 skipping at a concentration of 50 nM.

TABLE 38

(SEQ ID NOS 186-190, respectively, in order of appearance)				
Antisense oligonucleotide name	Sequence	Ability to induce skipping		
H52A(-07 + 14)	UCC UGC AUU GUU GCC UGU AAG	No skipping		
H52A(+12 + 41)	UCC AAC UGG GGA CGC CUC UGU UCC AAA UCC	Very strong skipping		

56

TABLE 38-continued

(SEQ ID NOS 186-190, respectively, in order of appearance)				
Antisense oligonucleotide name	Sequence	Ability to induce skipping		
H52A(+17 + 37)	ACU GGG GAC GCC UCU GUU CCA	Skipping to 50 nM		
H52A(+93 + 112)	CCG UAA UGA UUG UUC UAG CC	No skipping		
H52D(+05 - 15)	UGU UAA AAA ACU UAC UUC GA	No skipping		

Table 38 (SEQ ID NOS 186-190, respectively, in order of appearance)

Antisense Oligonucleotides Directed at Exon 53

Antisense oligonucleotides directed at exon 53 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

FIG. 22 also shows antisense molecule H53A(+39+69) [SEQ ID NO:193] directed at exon 53 acceptor splice site. This antisense oligonucleotide was able to induce exon 53 skipping at 5, 100, 300 and 600 nM. A "cocktail" of three exon 53 antisense oligonucleotides: H53A(+23+47) [SEQ ID NO:195], H53A(+150+176) [SEQ ID NO:196] and H53D(+14-07) [SEQ ID NO:194], was also tested, as shown in FIG. 20 and exhibited an ability to induce exon skipping.

Table 39 below includes other antisense molecules tested at a concentration range of 50, 100, 300 and 600 nM. These antisense molecules showed varying ability to induce exon 53 skipping. Antisense molecule H53A(+39+69) [SEQ ID NO:193] induced the strongest exon 53 skipping.

TABLE 39

(SEQ ID NOS 191-202, respectively, in order of appearance)				
Antisense oligonucleotide name	Sequence	Ability to induce skipping		
H53A(+45 + 69)	CAU UCA ACU GUU GCC UCC GGU UCU G	Faint skipping at 50 nM		
H53A(+39 + 62)	CUG UUG CCU CCG GUU CUG AAG GUG	Faint skipping at 50 nM		
H53A(+39 + 69)	CAU UCA ACU GUU GCC UCC GGU UCU GAA GGU G	Strong skipping to 50 nM		
H53D(+14 - 07)	UAC UAA CCU UGG UUU CUG UGA	Very faint skipping to 50 nM		
H53A(+23 + 47)	CUG AAG GUG UUC UUG UAC UUC AUC C	Very faint skipping to 50 nM		
H53A(+150 + 176)	UGU AUA GGG ACC CUC CUU CCA UGA CUC	Very faint skipping to 50 nM		
H53D(+20 - 05)	CUA ACC UUG GUU UCU GUG AUU UUC U	Not made yet		

US 8,455,636 B2

57

TABLE 39-continued

(SEQ ID NOS 191-202, respectively, in order of appearance)		
Antisense oligonucleotide name	Sequence	Ability to induce skipping
H53D(+09 - 18)	GGU AUC UUU GAU ACU AAC CUU GGU UUC	Faint at 600 nM
H53A(-12 + 10)	AUU CUU UCA ACU AGA AUA AAA G	No skipping
H53A(-07 + 18)	GAU UCU GAA UUC UUU CAA CUA GAA U	No skipping

58

TABLE 39-continued

(SEQ ID NOS 191-202, respectively, in order of appearance)		
Antisense oligonucleotide name	Sequence	Ability to induce skipping
H53A(+07 + 26)	AUC CCA CUG AUU CUG AAU UC	No skipping
H53A(+124 + 145)	UUG GCU CUG GCC UGU CCU AAG A	No skipping
15	Table 39 (SEQ ID NOS 191-202, respectively, in order of appearance)	

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 214

<210> SEQ ID NO 1
 <211> LENGTH: 24
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense
 oligonucleotide

<400> SEQUENCE: 1

gauagguggu aucaacaucu guaa

24

<210> SEQ ID NO 2
 <211> LENGTH: 21
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense
 oligonucleotide

<400> SEQUENCE: 2

gauagguggu aucaacaucu g

21

<210> SEQ ID NO 3
 <211> LENGTH: 25
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense
 oligonucleotide

<400> SEQUENCE: 3

gauagguggu aucaacaucu guaag

25

<210> SEQ ID NO 4
 <211> LENGTH: 20
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense
 oligonucleotide

<400> SEQUENCE: 4

US 8,455,636 B2

59

60

-continued

ggugguauca acaucuguaa 20

<210> SEQ ID NO 5
 <211> LENGTH: 20
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense
 oligonucleotide

<400> SEQUENCE: 5

guaucacau cuguaagcac 20

<210> SEQ ID NO 6
 <211> LENGTH: 23
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense
 oligonucleotide

<400> SEQUENCE: 6

ugcauguucc agucguugug ugg 23

<210> SEQ ID NO 7
 <211> LENGTH: 25
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense
 oligonucleotide

<400> SEQUENCE: 7

cacuaaucca gucaaaauagg ucugg 25

<210> SEQ ID NO 8
 <211> LENGTH: 25
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense
 oligonucleotide

<400> SEQUENCE: 8

auuuaccaac cuucaggauc gagua 25

<210> SEQ ID NO 9
 <211> LENGTH: 21
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense
 oligonucleotide

<400> SEQUENCE: 9

ggccuaaaac acauacacau a 21

<210> SEQ ID NO 10
 <211> LENGTH: 20
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Canine 2'-O-methyl phosphorothioate antisense

US 8,455,636 B2

61

62

-continued

oligonucleotide

<400> SEQUENCE: 10

cauuuuugac cuacaugugg 20

<210> SEQ ID NO 11
 <211> LENGTH: 20
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Canine 2'-O-methyl phosphorothioate antisense
 oligonucleotide

<400> SEQUENCE: 11

uuugaccuac auguggaaag 20

<210> SEQ ID NO 12
 <211> LENGTH: 26
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Canine 2'-O-methyl phosphorothioate antisense
 oligonucleotide

<400> SEQUENCE: 12

uacauuuuug accuacaugu ggaaag 26

<210> SEQ ID NO 13
 <211> LENGTH: 22
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Canine 2'-O-methyl phosphorothioate antisense
 oligonucleotide

<400> SEQUENCE: 13

auuuuugacc uacaugggaa ag 22

<210> SEQ ID NO 14
 <211> LENGTH: 23
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Canine 2'-O-methyl phosphorothioate antisense
 oligonucleotide

<400> SEQUENCE: 14

uacgaguuga uugucggacc cag 23

<210> SEQ ID NO 15
 <211> LENGTH: 25
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Canine 2'-O-methyl phosphorothioate antisense
 oligonucleotide

<400> SEQUENCE: 15

guggucuccu uaccuaugac ugugg 25

<210> SEQ ID NO 16
 <211> LENGTH: 17
 <212> TYPE: RNA

US 8,455,636 B2

63

64

-continued

```

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Canine 2'-O-methyl phosphorothioate antisense
      oligonucleotide

<400> SEQUENCE: 16

ggucuccuaa ccuauga                                17

<210> SEQ ID NO 17
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Human 2'-O-methyl phosphorothioate antisense
      oligonucleotide

<400> SEQUENCE: 17

ugucucagua aucuucuuac cuau                        24

<210> SEQ ID NO 18
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Human 2'-O-methyl phosphorothioate antisense
      oligonucleotide

<400> SEQUENCE: 18

ucuuaccuau gacuauggau gaga                        24

<210> SEQ ID NO 19
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Human 2'-O-methyl phosphorothioate antisense
      oligonucleotide

<400> SEQUENCE: 19

gcaugaacuc uuguggaucc                            20

<210> SEQ ID NO 20
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Human 2'-O-methyl phosphorothioate antisense
      oligonucleotide

<400> SEQUENCE: 20

ccagguacu acuuacauua                            20

<210> SEQ ID NO 21
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Human 2'-O-methyl phosphorothioate antisense
      oligonucleotide

<400> SEQUENCE: 21

aucguguguc acagcaucca g                          21

```

US 8,455,636 B2

65

66

-continued

<210> SEQ ID NO 22
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 22

uguuucagggc augaacucuu guggauccuu 30

<210> SEQ ID NO 23
<211> LENGTH: 31
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 23

uaggaggcgc cucccauccu guaggucacu g 31

<210> SEQ ID NO 24
<211> LENGTH: 31
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 24

aggucuagga ggcgccuccc auccguagg u 31

<210> SEQ ID NO 25
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 25

gcgccuccca uccuguaggu cacug 25

<210> SEQ ID NO 26
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 26

cuucgaggag gucuaggagg cgccuc 26

<210> SEQ ID NO 27
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

US 8,455,636 B2

67

68

-continued

<400> SEQUENCE: 27

cucccauccu guaggucacu g

21

<210> SEQ ID NO 28

<211> LENGTH: 22

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 28

uaccaguuuu ugcccuguca gg

22

<210> SEQ ID NO 29

<211> LENGTH: 26

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 29

ucaauaugcu gcuucccaaa cugaaa

26

<210> SEQ ID NO 30

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 30

cuaggaggcg ccucccaucc uguag

25

<210> SEQ ID NO 31

<211> LENGTH: 31

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 31

uuaugauuuc caucuacgau gucaguacuu c

31

<210> SEQ ID NO 32

<211> LENGTH: 31

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 32

cuuaccugcc aguggaggau uauauuccaa a

31

<210> SEQ ID NO 33

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

US 8,455,636 B2

69

70

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 33

caucaggauu cuuaccugcc agugg 25

<210> SEQ ID NO 34

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 34

cgaugucagu acuuccaaua uucac 25

<210> SEQ ID NO 35

<211> LENGTH: 18

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 35

accauucauc aggauucu 18

<210> SEQ ID NO 36

<211> LENGTH: 18

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 36

accugccagu ggaggauu 18

<210> SEQ ID NO 37

<211> LENGTH: 27

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 37

ccaauauuca cuaaaauaac cuguuuaa 27

<210> SEQ ID NO 38

<211> LENGTH: 30

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 38

caggauucu accugccagu ggaggauuau 30

<210> SEQ ID NO 39

US 8,455,636 B2

71

72

-continued

<211> LENGTH: 31
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 39

acgaugucag uacuuccaau auucacuaaa u 31

<210> SEQ ID NO 40
<211> LENGTH: 31
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 40

auuuccaucu acgaugucag uacuuccaau a 31

<210> SEQ ID NO 41
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 41

caggagcuuc caaavgcugc a 21

<210> SEQ ID NO 42
<211> LENGTH: 29
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 42

cuugucuca ggagcuucca aavgcugca 29

<210> SEQ ID NO 43
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 43

uccucagcag aaagaagcca cg 22

<210> SEQ ID NO 44
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 44

US 8,455,636 B2

73

74

-continued

uuagaaaucu cuccuugugc 20

<210> SEQ ID NO 45
 <211> LENGTH: 20
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense
 oligonucleotide
 <400> SEQUENCE: 45

uaaaauugggu guuacacaau 20

<210> SEQ ID NO 46
 <211> LENGTH: 24
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense
 oligonucleotide
 <400> SEQUENCE: 46

cccugaggca uucccaucuu gaau 24

<210> SEQ ID NO 47
 <211> LENGTH: 20
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense
 oligonucleotide
 <400> SEQUENCE: 47

aggacuuacu ugcuuuguuu 20

<210> SEQ ID NO 48
 <211> LENGTH: 23
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense
 oligonucleotide
 <400> SEQUENCE: 48

cuugaaauua ggagauucau cug 23

<210> SEQ ID NO 49
 <211> LENGTH: 23
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense
 oligonucleotide
 <400> SEQUENCE: 49

caucuucuga uaaauuuccu guu 23

<210> SEQ ID NO 50
 <211> LENGTH: 24
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense

US 8,455,636 B2

75

76

-continued

oligonucleotide

<400> SEQUENCE: 50

ucuucuguuu uuguuagcca guca 24

<210> SEQ ID NO 51

<211> LENGTH: 20

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 51

ucuauguaaa cugaaaauuu 20

<210> SEQ ID NO 52

<211> LENGTH: 20

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 52

uucuggagau ccauuaaaac 20

<210> SEQ ID NO 53

<211> LENGTH: 24

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 53

cagcaguugc gugaucucca cuag 24

<210> SEQ ID NO 54

<211> LENGTH: 21

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 54

uucaucaacu accaccacca u 21

<210> SEQ ID NO 55

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 55

cuaagcaaaa uaaucugacc uuaag 25

<210> SEQ ID NO 56

<211> LENGTH: 28

<212> TYPE: RNA

US 8,455,636 B2

77

78

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 56

cuuguaaaag aacccagcgg ucuucugu 28

<210> SEQ ID NO 57
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 57

caucuacaga uguuugccca uc 22

<210> SEQ ID NO 58
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 58

gaaggauguc uuguuaaaga acc 23

<210> SEQ ID NO 59
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 59

accguucuu caguaagacg 20

<210> SEQ ID NO 60
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 60

caugacacac cuguucuca guaa 24

<210> SEQ ID NO 61
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 61

cauuugagaa ggaugucuug 20

US 8,455,636 B2

79

80

-continued

<210> SEQ ID NO 62
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 62

aucucccaau accuggagaa gaga 24

<210> SEQ ID NO 63
<211> LENGTH: 31
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 63

gccaugcacu aaaaaggcac ugcaagacau u 31

<210> SEQ ID NO 64
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 64

ucuuuaaagc caguugugug aauc 24

<210> SEQ ID NO 65
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 65

uuucugaaag ccaugcacua a 21

<210> SEQ ID NO 66
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 66

guacauacgg ccaguuuuug aagac 25

<210> SEQ ID NO 67
<211> LENGTH: 31
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

US 8,455,636 B2

81

82

-continued

<400> SEQUENCE: 67

cuagaucgc uuuuaaaacc uguuaaaaca a

31

<210> SEQ ID NO 68

<211> LENGTH: 31

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 68

ucuuuucua g auccgcuuuu aaaaccuguu a

31

<210> SEQ ID NO 69

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 69

cuagaucgc uuuuaaaacc uguua

25

<210> SEQ ID NO 70

<211> LENGTH: 23

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 70

ccgucuuucg ggucacugac uua

23

<210> SEQ ID NO 71

<211> LENGTH: 26

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 71

cuagaucgc uuuuaaaacc uguuaa

26

<210> SEQ ID NO 72

<211> LENGTH: 20

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 72

ccgcuuuuaa aaccuguuaa

20

<210> SEQ ID NO 73

<211> LENGTH: 26

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

US 8,455,636 B2

83

84

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 73

uggauugcuu uuucuuuucu agaucc 26

<210> SEQ ID NO 74

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 74

caugcuuccg ucuucugggu cacug 25

<210> SEQ ID NO 75

<211> LENGTH: 23

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 75

gaucuuguuu gagugaauc agu 23

<210> SEQ ID NO 76

<211> LENGTH: 22

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 76

guuauccagc caugcuuccg uc 22

<210> SEQ ID NO 77

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 77

ugauaaauugg uaucacuaac cugug 25

<210> SEQ ID NO 78

<211> LENGTH: 22

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 78

guaucacuaa ccugugcugu ac 22

<210> SEQ ID NO 79

US 8,455,636 B2

85

86

-continued

<211> LENGTH: 19
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 79

cugcuggcau cuugcaguu 19

<210> SEQ ID NO 80
<211> LENGTH: 31
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 80

gccugagcug aucugcuggc aucuugcagu u 31

<210> SEQ ID NO 81
<211> LENGTH: 28
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 81

cuggcagaau ucgauccacc ggcuguuc 28

<210> SEQ ID NO 82
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 82

cagcaguagu ugucaucugc uc 22

<210> SEQ ID NO 83
<211> LENGTH: 19
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 83

ugauggggug guggguugg 19

<210> SEQ ID NO 84
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 84

US 8,455,636 B2

87

88

-continued

aucugcaua acaccucua gaaag 25

<210> SEQ ID NO 85
 <211> LENGTH: 24
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense
 oligonucleotide
 <400> SEQUENCE: 85

ccggcuguuc aguuguucug aggc 24

<210> SEQ ID NO 86
 <211> LENGTH: 28
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense
 oligonucleotide
 <400> SEQUENCE: 86

aucugcaua acaccucua gaaagaaa 28

<210> SEQ ID NO 87
 <211> LENGTH: 28
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense
 oligonucleotide
 <400> SEQUENCE: 87

gaaggagaag agauucuac cuuacaaa 28

<210> SEQ ID NO 88
 <211> LENGTH: 20
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense
 oligonucleotide
 <400> SEQUENCE: 88

auucgaucca ccggcuguuc 20

<210> SEQ ID NO 89
 <211> LENGTH: 20
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense
 oligonucleotide
 <400> SEQUENCE: 89

cagcaguagu ugucaucugc 20

<210> SEQ ID NO 90
 <211> LENGTH: 22
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense

US 8,455,636 B2

89

90

-continued

oligonucleotide

<400> SEQUENCE: 90

gccgguugac uucauccugu gc 22

<210> SEQ ID NO 91

<211> LENGTH: 22

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 91

cugcauccag gaacaugggu cc 22

<210> SEQ ID NO 92

<211> LENGTH: 23

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 92

gucugcaucc aggaacaugg guc 23

<210> SEQ ID NO 93

<211> LENGTH: 24

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 93

guugaagauc ugauagccgg uuga 24

<210> SEQ ID NO 94

<211> LENGTH: 24

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 94

uacuuacugu cuguagcucu uuuc 24

<210> SEQ ID NO 95

<211> LENGTH: 24

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 95

cacucauggu cuccugauag cgca 24

<210> SEQ ID NO 96

<211> LENGTH: 22

<212> TYPE: RNA

US 8,455,636 B2

91

92

-continued

```

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Human 2'-O-methyl phosphorothioate antisense
      oligonucleotide

<400> SEQUENCE: 96

cugcaauucc ccgagucucu gc                                22

<210> SEQ ID NO 97
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Human 2'-O-methyl phosphorothioate antisense
      oligonucleotide

<400> SEQUENCE: 97

acugcuggac ccauguccug aug                                23

<210> SEQ ID NO 98
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Human 2'-O-methyl phosphorothioate antisense
      oligonucleotide

<400> SEQUENCE: 98

cuaaguugag guauggagag u                                  21

<210> SEQ ID NO 99
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Human 2'-O-methyl phosphorothioate antisense
      oligonucleotide

<400> SEQUENCE: 99

uauucacaga ccugcaauuc ccc                                23

<210> SEQ ID NO 100
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Human 2'-O-methyl phosphorothioate antisense
      oligonucleotide

<400> SEQUENCE: 100

acaguggugc ugagauagua uaggcc                             26

<210> SEQ ID NO 101
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Human 2'-O-methyl phosphorothioate antisense
      oligonucleotide

<400> SEQUENCE: 101

uaggccacuu uguugcucuu gc                                22

```


US 8,455,636 B2

93

94

-continued

<210> SEQ ID NO 102
<211> LENGTH: 19
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 102

uucagagggc gcuuucuuc 19

<210> SEQ ID NO 103
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 103

gggcaggcca uuccuccuuc aga 23

<210> SEQ ID NO 104
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 104

ucuucagggg uuguauguga uucu 24

<210> SEQ ID NO 105
<211> LENGTH: 27
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 105

cugggcugaa uugucugaau aucacug 27

<210> SEQ ID NO 106
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 106

cuguuggcac augugaucac acugag 26

<210> SEQ ID NO 107
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

US 8,455,636 B2

95

96

-continued

<400> SEQUENCE: 107

gucuauaccu guuggcacau guga

24

<210> SEQ ID NO 108

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 108

ugcuuucugu aaaucaucug gaguu

25

<210> SEQ ID NO 109

<211> LENGTH: 26

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 109

ccuccuuucu ggcauagacc uuccac

26

<210> SEQ ID NO 110

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 110

ugugucaucc auucgugcau cucug

25

<210> SEQ ID NO 111

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 111

uuaaggccuc uuugucuaca ggugg

25

<210> SEQ ID NO 112

<211> LENGTH: 23

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 112

gggccucuuc uuuagcucuc uga

23

<210> SEQ ID NO 113

<211> LENGTH: 22

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

US 8,455,636 B2

97

98

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 113

gacuucaaaa gucuugcauu uc 22

<210> SEQ ID NO 114

<211> LENGTH: 24

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 114

gccaaacaugc ccaaacuucc uaa 24

<210> SEQ ID NO 115

<211> LENGTH: 26

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 115

cagagauuuc cucagcuccg ccagga 26

<210> SEQ ID NO 116

<211> LENGTH: 21

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 116

cuuacaucua gcaccucaga g 21

<210> SEQ ID NO 117

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 117

uccgccaucu guuagggucu gugcc 25

<210> SEQ ID NO 118

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 118

auuuggguua uccucugaau gucgc 25

<210> SEQ ID NO 119

US 8,455,636 B2

99

100

-continued

```

<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        Human 2'-O-methyl phosphorothioate antisense
        oligonucleotide

<400> SEQUENCE: 119

cauaccucuu cauguaguuc cc                                22


<210> SEQ ID NO 120
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        Human 2'-O-methyl phosphorothioate antisense
        oligonucleotide

<400> SEQUENCE: 120

cauuugagcu gcguccaccu ugucug                                26


<210> SEQ ID NO 121
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        Human 2'-O-methyl phosphorothioate antisense
        oligonucleotide

<400> SEQUENCE: 121

uccugggcag acuggaugcu cuguuc                                26


<210> SEQ ID NO 122
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        Human 2'-O-methyl phosphorothioate antisense
        oligonucleotide

<400> SEQUENCE: 122

uugccugggc uuccugaggc auu                                23


<210> SEQ ID NO 123
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        Human 2'-O-methyl phosphorothioate antisense
        oligonucleotide

<400> SEQUENCE: 123

uucugaaaau acauauaccu gugc                                24


<210> SEQ ID NO 124
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        Human 2'-O-methyl phosphorothioate antisense
        oligonucleotide

<400> SEQUENCE: 124

```

US 8,455,636 B2

101

102

-continued

uaguuucuga aauaacaauau accug 25

<210> SEQ ID NO 125
 <211> LENGTH: 21
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense
 oligonucleotide

<400> SEQUENCE: 125

gacuugucua aucagauugg a 21

<210> SEQ ID NO 126
 <211> LENGTH: 24
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense
 oligonucleotide

<400> SEQUENCE: 126

guuucugaaa uaacauauac cugu 24

<210> SEQ ID NO 127
 <211> LENGTH: 20
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense
 oligonucleotide

<400> SEQUENCE: 127

caccagaaau acauaccaca 20

<210> SEQ ID NO 128
 <211> LENGTH: 20
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense
 oligonucleotide

<400> SEQUENCE: 128

caaugauuuu gcugugacug 20

<210> SEQ ID NO 129
 <211> LENGTH: 23
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense
 oligonucleotide

<400> SEQUENCE: 129

cgaaacuua uggagacau uug 23

<210> SEQ ID NO 130
 <211> LENGTH: 25
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense

US 8,455,636 B2

103

104

-continued

oligonucleotide

<400> SEQUENCE: 130

cuuguagacg cugcucaaaa uuggc 25

<210> SEQ ID NO 131

<211> LENGTH: 20

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 131

caugcacaca ccuuugcucc 20

<210> SEQ ID NO 132

<211> LENGTH: 24

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 132

ucuguacaau cugacgucca gucu 24

<210> SEQ ID NO 133

<211> LENGTH: 27

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 133

gucuuuauca ccauuuccac uucagac 27

<210> SEQ ID NO 134

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 134

ccgucugcuu uuucuguaca aucug 25

<210> SEQ ID NO 135

<211> LENGTH: 22

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 135

uccauaucug uagcugccag cc 22

<210> SEQ ID NO 136

<211> LENGTH: 23

<212> TYPE: RNA

US 8,455,636 B2

105

106

-continued

```

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Human 2'-O-methyl phosphorothioate antisense
      oligonucleotide

<400> SEQUENCE: 136

ccaggcaacu ucagaaucca aau                                     23

<210> SEQ ID NO 137
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Human 2'-O-methyl phosphorothioate antisense
      oligonucleotide

<400> SEQUENCE: 137

uuucuguuac cugaaaagaa uuauaaugaa                             30

<210> SEQ ID NO 138
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Human 2'-O-methyl phosphorothioate antisense
      oligonucleotide

<400> SEQUENCE: 138

cauucuuuuc cuuucgcauc uuacg                                   25

<210> SEQ ID NO 139
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Human 2'-O-methyl phosphorothioate antisense
      oligonucleotide

<400> SEQUENCE: 139

ugaucucuuu gucaauucca uaucug                                   26

<210> SEQ ID NO 140
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Human 2'-O-methyl phosphorothioate antisense
      oligonucleotide

<400> SEQUENCE: 140

uucagugaua uagguuuuac cuuuccccag                             30

<210> SEQ ID NO 141
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Human 2'-O-methyl phosphorothioate antisense
      oligonucleotide

<400> SEQUENCE: 141

cuguagcugc cagccauucu gucaag                                 26

```

US 8,455,636 B2

107

108

-continued

<210> SEQ ID NO 142
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 142

ucuucugcuc gggaggugac a 21

<210> SEQ ID NO 143
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 143

ccaguuacua uucagaagac 20

<210> SEQ ID NO 144
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 144

ucuucaggug caccuucugu 20

<210> SEQ ID NO 145
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 145

ugugaugugg uccacauucu gguca 25

<210> SEQ ID NO 146
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 146

ccauguguuu cugguauucc 20

<210> SEQ ID NO 147
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

US 8,455,636 B2

109

110

-continued

<400> SEQUENCE: 147

cguguagagu ccaccuuugg gcgua

25

<210> SEQ ID NO 148

<211> LENGTH: 24

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 148

uacuaauuuc cugcaguggu cacc

24

<210> SEQ ID NO 149

<211> LENGTH: 24

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 149

uucuguguga aauggcugca aauc

24

<210> SEQ ID NO 150

<211> LENGTH: 20

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 150

ccuucuaaagg aauggaggcc

20

<210> SEQ ID NO 151

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 151

ugcugaauuu cagccuccag ugguu

25

<210> SEQ ID NO 152

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 152

ugaagucuuc cucuuucaga uucac

25

<210> SEQ ID NO 153

<211> LENGTH: 24

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

US 8,455,636 B2

111

112

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 153

cuggcuuucu cucaucugug auuc 24

<210> SEQ ID NO 154

<211> LENGTH: 20

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense oligonucleotide

<400> SEQUENCE: 154

guuguaaguu guccuccuu 20

<210> SEQ ID NO 155

<211> LENGTH: 20

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 155

uugucuguaa cagcugcugu 20

<210> SEQ ID NO 156

<211> LENGTH: 20

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 156

gcucuaauac cuugagagca 20

<210> SEQ ID NO 157

<211> LENGTH: 22

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 157

cuuugagacc ucaaauccug uu 22

<210> SEQ ID NO 158

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 158

cuuuuuuuuc cuuucaucuc ugggc 25

<210> SEQ ID NO 159

<211> LENGTH: 27

US 8,455,636 B2

113

114

-continued

<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 159

aucguuucuu cacggacagu gugcugg 27

<210> SEQ ID NO 160
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 160

gggcuuguga gacaugagug auuu 24

<210> SEQ ID NO 161
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 161

accuucagag gacuccucuu gc 22

<210> SEQ ID NO 162
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 162

uauuguuac cuacccuugu cgguc 25

<210> SEQ ID NO 163
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 163

ggagagagcu uccuguagcu 20

<210> SEQ ID NO 164
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 164

ucacccuuc cacaggcguu gca 23

US 8,455,636 B2

115

116

-continued

<210> SEQ ID NO 165
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 165

uuugugucuu ucugagaaac 20

<210> SEQ ID NO 166
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 166

aaagacuuac cuuaagauac 20

<210> SEQ ID NO 167
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 167

aucugucaaa ucgccugcag 20

<210> SEQ ID NO 168
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 168

uuaccuugac uugcucaagc 20

<210> SEQ ID NO 169
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 169

uccagguuca agugggauac 20

<210> SEQ ID NO 170
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

US 8,455,636 B2

117

118

-continued

<400> SEQUENCE: 170

gcucuucugg gcuuauggga gcacu

25

<210> SEQ ID NO 171

<211> LENGTH: 27

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 171

accuuuaucc acuggagauu ugucugc

27

<210> SEQ ID NO 172

<211> LENGTH: 21

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 172

uuccaccagu aacugaaaca g

21

<210> SEQ ID NO 173

<211> LENGTH: 29

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 173

ccacucagag cucagauuu cuaacuucc

29

<210> SEQ ID NO 174

<211> LENGTH: 27

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 174

cuuccacuca gagcucagau cuucuaa

27

<210> SEQ ID NO 175

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 175

gggauccagu auacuuacag gcucc

25

<210> SEQ ID NO 176

<211> LENGTH: 26

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

US 8,455,636 B2

119

120

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 176

accagaguua cagucugagu aggagc 26

<210> SEQ ID NO 177
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 177

cucauaccuu cugcuugaug auc 23

<210> SEQ ID NO 178
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 178

uucuguccaa gcccgguuga aauc 24

<210> SEQ ID NO 179
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 179

acaucaagga agauggcauu ucuaguugg 30

<210> SEQ ID NO 180
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 180

acaucaagga agauggcauu ucuag 25

<210> SEQ ID NO 181
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 181

cuccaacauc aaggaagaug gcuuuucuag 30

US 8,455,636 B2

121

122

-continued

<210> SEQ ID NO 182
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 182

aucauuuuuu cucauaccuu cugcu 25

<210> SEQ ID NO 183
<211> LENGTH: 36
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 183

aucauuuuuu cucauaccuu cugcuaggag cuaaaa 36

<210> SEQ ID NO 184
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 184

cacccaccau caccucugu g 21

<210> SEQ ID NO 185
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 185

aucaucucgu ugauauccuc aa 22

<210> SEQ ID NO 186
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 186

uccugcaug uugccguaa g 21

<210> SEQ ID NO 187
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 187

US 8,455,636 B2

123

124

-continued

uccaacuggg gacgccucug uuccaaaucc 30

<210> SEQ ID NO 188
 <211> LENGTH: 21
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense
 oligonucleotide

<400> SEQUENCE: 188

acuggggacg ccucuguucc a 21

<210> SEQ ID NO 189
 <211> LENGTH: 20
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense
 oligonucleotide

<400> SEQUENCE: 189

ccguaaagau uguucuagcc 20

<210> SEQ ID NO 190
 <211> LENGTH: 20
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense
 oligonucleotide

<400> SEQUENCE: 190

uguuaaaaaa cuuacuucga 20

<210> SEQ ID NO 191
 <211> LENGTH: 25
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense
 oligonucleotide

<400> SEQUENCE: 191

cauuaacug uugccuccgg uucug 25

<210> SEQ ID NO 192
 <211> LENGTH: 24
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Human 2'-O-methyl phosphorothioate antisense
 oligonucleotide

<400> SEQUENCE: 192

cuguugccuc cgguucugaa ggug 24

<210> SEQ ID NO 193
 <211> LENGTH: 31
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

US 8,455,636 B2

125

126

-continued

Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 193

cauucacacug uugccuccgg uucugaaggu g 31

<210> SEQ ID NO 194
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 194

uacuaaccuu gguuucugug a 21

<210> SEQ ID NO 195
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 195

cugaaggugu ucuuguacuu caucc 25

<210> SEQ ID NO 196
<211> LENGTH: 27
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 196

uguauaggga cccuccuucc augacuc 27

<210> SEQ ID NO 197
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 197

cuaaccuugg uuucugugau uuucu 25

<210> SEQ ID NO 198
<211> LENGTH: 27
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 198

gguaucuuug auacuaaccu ugguuuc 27

<210> SEQ ID NO 199
<211> LENGTH: 22

US 8,455,636 B2

127

128

-continued

```

<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Human 2'-O-methyl phosphorothioate antisense
      oligonucleotide

<400> SEQUENCE: 199

auucuuucaa cuagaauaaa ag                                22

<210> SEQ ID NO 200
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Human 2'-O-methyl phosphorothioate antisense
      oligonucleotide

<400> SEQUENCE: 200

gauucugaau ucuuuaacu agaau                                25

<210> SEQ ID NO 201
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Human 2'-O-methyl phosphorothioate antisense
      oligonucleotide

<400> SEQUENCE: 201

aucccacuga uucugaaauuc                                    20

<210> SEQ ID NO 202
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Human 2'-O-methyl phosphorothioate antisense
      oligonucleotide

<400> SEQUENCE: 202

uuggcucugg ccuguccuaa ga                                22

<210> SEQ ID NO 203
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Human 2'-O-methyl phosphorothioate antisense
      oligonucleotide

<400> SEQUENCE: 203

cucuuuuucca gguucaagug ggauacuagc                        30

<210> SEQ ID NO 204
<211> LENGTH: 31
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Human 2'-O-methyl phosphorothioate antisense
      oligonucleotide

<400> SEQUENCE: 204

caagcuuuuc uuuuaguugc ugcucuuuuc c                        31

```


US 8,455,636 B2

129

130

-continued

<210> SEQ ID NO 205
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 205

uauucuuuug uucuucuaag cuggagaaag 30

<210> SEQ ID NO 206
<211> LENGTH: 28
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 206

cugcuuccuc caaccauaaa acaaaauuc 28

<210> SEQ ID NO 207
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 207

ccaaugccau ccuggaguuc cuguuaa 26

<210> SEQ ID NO 208
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 208

uccuguagaa uacuggcauc 20

<210> SEQ ID NO 209
<211> LENGTH: 27
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 209

ugcagaccuc cugccaccgc agauuca 27

<210> SEQ ID NO 210
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

US 8,455,636 B2

131

132

-continued

<400> SEQUENCE: 210

cuaccucuuu uuucugucug

20

<210> SEQ ID NO 211

<211> LENGTH: 20

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 211

uguuuuugag gauugcugaa

20

<210> SEQ ID NO 212

<211> LENGTH: 84

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 212

cagcaguagu ugucaucugc ucaacuggca gaauucgauc caccggcugu ucaagccuga

60

gcugaucugc ucgcaucuug cagu

84

<210> SEQ ID NO 213

<211> LENGTH: 44

<212> TYPE: RNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 213

uauagcacug agugaccucu uucucgcagg cgcuaugcugg agca

44

<210> SEQ ID NO 214

<211> LENGTH: 22

<212> TYPE: RNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 214

ccgugcagac ugacggucuc au

22

The claims defining the invention are as follows:

1. An isolated antisense oligonucleotide of 20 to 50 nucleotides in length comprising at least 20 consecutive nucleotides of SEQ ID NO:193, wherein the oligonucleotide specifically hybridizes to an exon 53 target region of the human dystrophin gene inducing exon 53 skipping, and wherein the uracil bases are optionally thymine bases.

2. The antisense oligonucleotide of claim 1 comprising SEQ ID NO:193.

3. The antisense oligonucleotide of claim 1 consisting of SEQ ID NO:193.

4. The antisense oligonucleotide of claim 1 comprising 20-31 nucleotides in length.

5. The antisense oligonucleotide of claim 1, wherein the oligonucleotide does not activate RNase H.

6. The antisense oligonucleotide of claim 1, comprising a non-natural backbone.

7. The antisense oligonucleotide of claim 1, wherein the sugar moieties of the oligonucleotide backbone are replaced with non-natural moieties.

8. The antisense oligonucleotide of claim 7, wherein the non-natural moieties are morpholinos.

9. The antisense oligonucleotide of claim 1, wherein the inter-nucleotide linkages of the oligonucleotide backbone are replaced with non-natural inter-nucleotide linkages.

10. The antisense oligonucleotide of claim 9, wherein the non-natural inter-nucleotide linkages are modified phosphates.

11. The antisense oligonucleotide of claim 1, wherein the sugar moieties of the oligonucleotide backbone are replaced with non-natural moieties and the inter-nucleotide linkages of the oligonucleotide backbone are replaced with non-natural inter-nucleotide linkages.

12. The antisense oligonucleotide of claim 11, wherein the non-natural moieties are morpholinos and the non-natural internucleotide linkages are modified phosphates.

13. The antisense oligonucleotide of claim 12, wherein the modified phosphates are methyl phosphonates, methyl phosphorothioates, phosphoromorpholidates, phosphoropiperazides or phosphoramidates.

US 8,455,636 B2

133

14. The antisense oligonucleotide of claim 1, wherein the oligonucleotide is a 2'-O-methyl-oligoribonucleotide.

15. The antisense oligonucleotide of claim 1, wherein the oligonucleotide is a peptide nucleic acid.

16. The antisense oligonucleotide of claim 1, wherein the oligonucleotide is chemically linked to one or more moieties or conjugates that enhance the activity, cellular distribution, or cellular uptake of the antisense oligonucleotide.

17. The antisense oligonucleotide of claim 16, wherein the oligonucleotide is conjugated to a polyamine.

18. The antisense oligonucleotide of claim 16, wherein the oligonucleotide is chemically linked to a polyethylene glycol chain.

19. An isolated antisense oligonucleotide of 20 to 50 nucleotides in length comprising at least 20 consecutive nucleotides complementary to an exon 53 target region of the human dystrophin gene designated as annealing site H53A(+39+69), wherein the antisense oligonucleotide specifically hybridizes to the annealing site inducing exon 53 skipping, and wherein uracil bases in the antisense oligonucleotide are optionally thymine bases.

20. The antisense oligonucleotide of claim 19 comprising 20-31 nucleotides in length.

21. The antisense oligonucleotide of claim 19, wherein the uracil bases are thymine.

22. The antisense oligonucleotide of claim 19, wherein the oligonucleotide does not activate RNase H.

23. The antisense oligonucleotide of claim 19, comprising a non-natural backbone.

24. The antisense oligonucleotide of claim 19, wherein the sugar moieties of the oligonucleotide backbone are replaced with non-natural moieties.

25. The antisense oligonucleotide of claim 24, wherein the non-natural moieties are morpholinos.

26. The antisense oligonucleotide of claim 19, wherein the inter-nucleotide linkages of the oligonucleotide backbone are replaced with non-natural inter-nucleotide linkages.

27. The antisense oligonucleotide of claim 26, wherein the non-natural inter-nucleotide linkages are modified phosphates.

28. The antisense oligonucleotide of claim 19, wherein the sugar moieties of the oligonucleotide backbone are replaced with non-natural moieties and the inter-nucleotide linkages of the oligonucleotide backbone are replaced with non-natural inter-nucleotide linkages.

134

29. The antisense oligonucleotide of claim 28, wherein the non-natural moieties are morpholinos and the non-natural internucleotide linkages are modified phosphates.

30. The antisense oligonucleotide of claim 29, wherein the modified phosphates are methyl phosphonates, methyl phosphorothioates, phosphoromorpholidates, phosphoropiperazides or phosphoroamidates.

31. The antisense oligonucleotide of claim 19, wherein the oligonucleotide is a 2'-O-methyl-oligoribonucleotide.

32. The antisense oligonucleotide of claim 19, wherein the oligonucleotide is a peptide nucleic acid.

33. The antisense oligonucleotide of claim 19, wherein the oligonucleotide is chemically linked to one or more moieties or conjugates that enhance the activity, cellular distribution, or cellular uptake of the antisense oligonucleotide.

34. The antisense oligonucleotide of claim 33, wherein the oligonucleotide is conjugated to a polyamine.

35. The antisense oligonucleotide of claim 33, wherein the oligonucleotide is chemically linked to a polyethylene glycol chain.

36. A pharmaceutical composition, comprising an antisense oligonucleotide of claim 1, and a saline solution that includes a phosphate buffer.

37. The antisense oligonucleotide of claim 1, wherein the uracil bases are thymine bases.

38. The antisense oligonucleotide of claim 1, comprising SEQ ID NO:193, wherein the uracil bases are thymine bases.

39. A pharmaceutical composition, comprising an antisense oligonucleotide of claim 2, and a saline solution that includes a phosphate buffer.

40. A pharmaceutical composition, comprising an antisense oligonucleotide of claim 3, and a saline solution that includes a phosphate buffer.

41. A pharmaceutical composition, comprising an antisense oligonucleotide of claim 19, and a saline solution that includes a phosphate buffer.

42. A pharmaceutical composition, comprising an antisense oligonucleotide of claim 38, and a saline solution that includes a phosphate buffer.

43. A method of treating Duchenne muscular dystrophy, comprising administering to a patient in need thereof an effective amount of a pharmaceutical composition of claim 36.

* * * * *